

DCCG'S NATIONALE RETNINGSLINIER FOR DIAGNOSTIK OG BEHANDLING AF KOLOREKTAL CANCER		
Litteraturliste - Tromboseprofylakse		
Forfattere: MSN	Gælder fra: 1. December 2012	Gælder til:

- 1) Mismetti P, Laporte S, Darmon JY, Buchmüller A, Decousus H. Meta-analysis of low molecular weight heparin in the prevention of venous thromboembolism in general surgery. *Br J Surg* . 2001; 88 (7): 913 - 930.
- 2) Kakkar, V. V., et al. Low molecular weight versus standard heparin for prevention of venous thromboembolism after major abdominal surgery. The Thromboprophylaxis Collaborative Group. *Lancet* 1993; 341: 259-65.
- 3) Enoxacan Study Group. Efficacy and safety of enoxaparin versus unfractionated heparin for prevention of deep vein thrombosis in elective cancer surgery: a double-blind randomised multicentre trial with venographic assessment. *Br J Surg* 1997; 84:1099-103.
- 4) Oster, G., R. L. Tuden, and G. A. Colditz. Prevention of venous thromboembolism after general surgery. Cost-effectiveness analysis of alternative approaches to prophylaxis. *Am J Med* 1987; 889-99.
- 5) Persson, A. V., R. J. Davis, and J. L. Villavicencio. Deep venous thrombosis and pulmonary embolism. *Surg Clin North Am* 1991; 71: 1195-209.
- 6) Rickles FR, Levine MN. Epidemiology of thrombosis in cancer. *Acta Haematol.* 2001; 106: 6-12.
- 7) Wille-Jørgensen P, Rasmussen MS, Andersen BR, Borly L. Heparins and mechanical methods for thromboprophylaxis in colorectal surgery. *Cochrane Database Syst Rev* 2003;CD001217.
- 8) Stender MT, Nielsen TS, Frokjaer JB, Larsen TB, Lundbye-Christensen S, Thorlacius-Ussing O. High preoperative prevalence of deep venous thrombosis in patients with colorectal cancer. *Br J Surg* 2007;94:1100-3.8
- 9) Dentali F, Ageno W, Becattini C et al. Prevalence and clinical history of incidental, asymptomatic pulmonary embolism: a meta-analysis. *Thromb Res.* 2010 Jun;125(6):518-22.
- 10) [Bergqvist D, Burmark US, Flordal PA, Frisell J, Hallbook T, Hedberg M et al. Low molecular weight heparin started before surgery as prophylaxis against deep vein thrombosis: 2500 versus 5000 Xal units in 2070 patients. *Br J Surg* 1995;82:496-501).
- 11) Wiig JN, Solhaug JH, Bilberg T, Bjerkeset T, Edwin B, Gruner OP et al. Prophylaxis of venographically diagnosed deep vein thrombosis in gastrointestinal surgery. Multicentre trials 20 mg and 40 mg enoxaparin versus dextran. *Eur J Surg* 1995;161:663-8).

DCCG'S NATIONALE RETNINGSLINIER FOR DIAGNOSTIK OG BEHANDLING AF KOLOREKTAL CANCER		
Litteraturliste - Tromboseprofylakse		
Forfattere: MSN	Gælder fra: 1. December 2012	Gælder til:

- 12) Retningslinjer for perioperativ regulering af antitrombotisk behandling. 2011. www.DTSH.dk
- 13) Geerts WH, Bergqvist D, Pineo GF, et al. Prevention of venous thromboembolism: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). Chest 2008;133:381S-453S.
- 14) Bergqvist D, Agnelli G, Cohen AT, et al; ENOXACAN II Investigators . Duration of prophylaxis against venous thromboembolism with enoxaparin after surgery for cancer . N Engl J Med. 2002; 346 (13): 975 - 980.
- 15) Lausen I, Jensen R, Jorgensen LN, Rasmussen MS, Lyng KM, Andersen M, Raaschou, Wille-Jørgensen P. Incidence and prevention of deep venous thrombosis occurring late after general surgery: Randomised controlled study of prolonged thromboprophylaxis. Eur J Surg 1998;164:657-663
- 16) Rasmussen MS , Jorgensen LN , Wille-Jorgensen P , et al . Prolonged prophylaxis with dalteparin to prevent late thromboembolic complications in patients undergoing major abdominal surgery: a multicenter randomized open-label study . J Thromb Haemost . 2006; 4 (11): 2384 - 2390 .
- 17) Kakkar VV, Balibrea JL, Martínez-González J, Prandoni P; CANBESURE Study Group . Extended prophylaxis with bemiparin for the prevention of venous thromboembolism after abdominal or pelvic surgery for cancer: the CANBESURE randomized study . J Thromb Haemost . 2010; 8 (6) : 1223 - 1229 .
- 18) Rasmussen MS, Jørgensen LN, Wille-Jørgensen P. Prolonged thromboprophylaxis with low molecular weight heparin for abdominal or pelvic surgery. Cochrane Database Syst Rev .2009;(1): CD004318.
- 19) Behandvejledning for tromboseprofylakse til parenchymkirurgiske patienter. www.regioner.dk/sundhed/medicin/RADS/behandlingsvejledninger.
- 20) Beebe DS, McMavin MP, Crain JM, et al. Evidence of venous stasis after abdominal insufflation for laparoscopic cholecystectomy. Surg Gynecol Obst 1993; 176: 443-7
- 21) Guidelines for deep venous thrombosis prophylaxis during laparoscopic surgery. Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) Guidelines Committee. Surg Endosc. 2007 Jun;21(6):1007-9. Epub 2007 Apr 5