

DCCG'S NATIONALE RETNINGSLINIER FOR DIAGNOSTIK OG BEHANDLING AF KOLOREKTAL CANCER		
Litteraturliste – MR-staging af rectumcancer		
Forfattere: BGP	Gælder fra: 1. oktober 2012	Gælder til:

Reference List

- (1) Brown G, Richards CJ, Newcombe RG, Dallimore NS, Radcliffe AG, Carey DP et al. Rectal carcinoma: thin-section MR imaging for staging in 28 patients. *Radiology* 1999; 211(1):215-222.
- (2) Brown G, Radcliffe AG, Newcombe RG, Dallimore NS, Bourne MW, Williams GT. Preoperative assessment of prognostic factors in rectal cancer using high-resolution magnetic resonance imaging. *Br J Surg* 2003; 90(3):355-364.
- (3) Brown G, Daniels IR, Richardson C, Revell P, Peppercorn D, Bourne M. Techniques and trouble-shooting in high spatial resolution thin slice MRI for rectal cancer. *Br J Radiol* 2005; 78(927):245-251.
- (4) Brown G, Kirkham A, Williams GT, Bourne M, Radcliffe AG, Sayman J et al. High-resolution MRI of the anatomy important in total mesorectal excision of the rectum. *AJR Am J Roentgenol* 2004; 182(2):431-439.
- (5) Pedersen BG, Blomqvist L, Brown G, Fenger-Gron M, Moran B, Laurberg S. Postgraduate multidisciplinary development program: impact on the interpretation of pelvic MRI in patients with rectal cancer: a clinical audit in West Denmark. *Dis Colon Rectum* 2011; 54(3):328-334.
- (6) Burton S, Brown G, Daniels IR, Norman AR, Mason B, Cunningham D. MRI directed multidisciplinary team preoperative treatment strategy: the way to eliminate positive circumferential margins? *Br J Cancer* 2006; 94(3):351-357.
- (7) Palmer G, Martling A, Cedermark B, Holm T. Preoperative tumour staging with multidisciplinary team assessment improves the outcome in locally advanced primary rectal cancer. *Colorectal Dis* 2011; 13(12):1361-1369.
- (8) Merkel S, Mansmann U, Siassi M, Papadopoulos T, Hohenberger W, Hermanek P. The prognostic inhomogeneity in pT3 rectal carcinomas. *Int J Colorectal Dis* 2001; 16(5):298-304.
- (9) Willett CG, Badizadegan K, Ancukiewicz M, Shellito PC. Prognostic factors in stage T3N0 rectal cancer: do all patients require postoperative pelvic irradiation and chemotherapy? *Dis Colon Rectum* 1999; 42(2):167-173.
- (10) Steel MC, Woods R, Mackay JM, Chen F. Extent of mesorectal invasion is a prognostic indicator in T3 rectal carcinoma. *ANZ J Surg* 2002; 72(7):483-487.
- (11) Miyoshi M, Ueno H, Hashiguchi Y, Mochizuki H, Talbot IC. Extent of mesorectal tumor invasion as a prognostic factor after curative surgery for T3 rectal cancer patients. *Ann Surg* 2006; 243(4):492-498.
- (12) Extramural depth of tumor invasion at thin-section MR in patients with rectal cancer: results of the MERCURY study. *Radiology* 2007; 243(1):132-139.
- (13) Pedersen BG, Moran B, Brown G, Blomqvist L, Fenger-Gron M, Laurberg S. Reproducibility of Depth of Extramural Tumor Spread and Distance to Circumferential Resection Margin at Rectal MRI:

Litteraturliste – MR-staging af rectumcancer

Forfattere: BGP

Gælder fra: 1. oktober 2012

Gælder til:

- Enhancement of Clinical Guidelines for Neoadjuvant Therapy. *AJR Am J Roentgenol* 2011; 197(6):1360-1366.
- (14) Beets-Tan RG, Beets GL, Vliegen RF, Kessels AG, Van Boven H, De Bruine A et al. Accuracy of magnetic resonance imaging in prediction of tumour-free resection margin in rectal cancer surgery. *Lancet* 2001; 357(9255):497-504.
- (15) Nagtegaal ID, Marijnen CA, Kranenbarg EK, van de Velde CJ, van Krieken JH. Circumferential margin involvement is still an important predictor of local recurrence in rectal carcinoma: not one millimeter but two millimeters is the limit. *Am J Surg Pathol* 2002; 26(3):350-357.
- (16) Wibe A, Rendedal PR, Svensson E, Norstein J, Eide TJ, Myrvold HE et al. Prognostic significance of the circumferential resection margin following total mesorectal excision for rectal cancer. *Br J Surg* 2002; 89(3):327-334.
- (17) Diagnostic accuracy of preoperative magnetic resonance imaging in predicting curative resection of rectal cancer: prospective observational study. *BMJ* 2006; 333(7572):779.
- (18) Taylor FG, Quirke P, Heald RJ, Moran B, Blomqvist L, Swift I et al. One millimetre is the safe cut-off for magnetic resonance imaging prediction of surgical margin status in rectal cancer. *Br J Surg* 2011; 98(6):872-879.
- (19) Blomqvist L, Glimelius B. The 'good', the 'bad', and the 'ugly' rectal cancers. *Acta Oncol* 2008; 47(1):5-8.
- (20) Smith N, Brown G. Preoperative staging of rectal cancer. *Acta Oncol* 2008; 47(1):20-31.
- (21) Lahaye MJ, Engelen SM, Nelemans PJ, Beets GL, van de Velde CJ, van Engelshoven JM et al. Imaging for predicting the risk factors--the circumferential resection margin and nodal disease--of local recurrence in rectal cancer: a meta-analysis. *Semin Ultrasound CT MR* 2005; 26(4):259-268.
- (22) Bipat S, Glas AS, Slors FJ, Zwinderman AH, Bossuyt PM, Stoker J. Rectal cancer: local staging and assessment of lymph node involvement with endoluminal US, CT, and MR imaging--a meta-analysis. *Radiology* 2004; 232(3):773-783.
- (23) Klerkx WM, Bax L, Veldhuis WB, Heintz AP, Mali WP, Peeters PH et al. Detection of lymph node metastases by gadolinium-enhanced magnetic resonance imaging: systematic review and meta-analysis. *J Natl Cancer Inst* 2010; 102(4):244-253.
- (24) Brown G, Richards CJ, Bourne MW, Newcombe RG, Radcliffe AG, Dallimore NS et al. Morphologic predictors of lymph node status in rectal cancer with use of high-spatial-resolution MR imaging with histopathologic comparison. *Radiology* 2003; 227(2):371-377.
- (25) Al Sukhni E, Milot L, Fruitman M, Beyene J, Victor JC, Schmocker S et al. Diagnostic Accuracy of MRI for Assessment of T Category, Lymph Node Metastases, and Circumferential Resection Margin Involvement in Patients with Rectal Cancer: A Systematic Review and Meta-analysis. *Ann Surg Oncol* 2012.
- (26) Koh DM, George C, Temple L, Collins DJ, Toomey P, Raja A et al. Diagnostic accuracy of nodal enhancement pattern of rectal cancer at

DCCG'S NATIONALE RETNINGSLINIER FOR DIAGNOSTIK OG BEHANDLING AF KOLOREKTAL CANCER		
Litteraturliste – MR-staging af rectumcancer		
Forfattere: BGP	Gælder fra: 1. oktober 2012	Gælder til:

- MRI enhanced with ultrasmall superparamagnetic iron oxide: findings in pathologically matched mesorectal lymph nodes. *AJR Am J Roentgenol* 2010; 194(6):W505-W513.
- (27) Lahaye MJ, Engelen SM, Kessels AG, de Bruine AP, von Meyenfeldt MF, van Engelshoven JM et al. USPIO-enhanced MR imaging for nodal staging in patients with primary rectal cancer: predictive criteria. *Radiology* 2008; 246(3):804-811.
- (28) Patel UB, Taylor F, Blomqvist L, George C, Evans H, Tekkis P et al. Magnetic resonance imaging-detected tumor response for locally advanced rectal cancer predicts survival outcomes: MERCURY experience. *J Clin Oncol* 2011; 29(28):3753-3760.
- (29) Maizlin ZV, Brown JA, So G, Brown C, Phang TP, Walker ML et al. Can CT replace MRI in preoperative assessment of the circumferential resection margin in rectal cancer? *Dis Colon Rectum* 2010; 53(3):308-314.
- (30) Vliegen R, Dresen R, Beets G, Daniels-Gooszen A, Kessels A, van Engelshoven J et al. The accuracy of Multi-detector row CT for the assessment of tumor invasion of the mesorectal fascia in primary rectal cancer. *Abdom Imaging* 2008; 33(5):604-610.