

LITTERATUR

Abdalla EK et al. Improving Resectability of Hepatic Colorectal Metastases: Expert Consensus Statement. *Ann Surg Oncol* 2006; 13: 1271-80.

Abdalla EK et al. Recurrence and outcomes following hepatic resection, radiofrequency ablation, and combined resection/ablation for colorectal liver metastases. *Ann Surg* 2004; 239: 818-25.

Abraham NS, Young JM, Solomon MJ. Meta-analysis of short-term outcomes after laparoscopic resection for colorectal cancer. *Br J Surg* 2004; 91: 1111-1124. Achiam 2008.

Agnelli G, Bergqvist D, Cohen A, Gallus AS, Gent M. Randomized clinical trial of postoperative fondaparinux versus perioperative dalteparin for prevention of venous thrombosis in high-risk abdominal surgery. *Br J Surg* 2005; 92: 1212-1220.

Akasu T, Yamaguchi T, Fujimoto Y, Ishiguro S, Yamamoto S, Fujita S et al. Abdominal sacral resection for posterior pelvic recurrence of rectal carcinoma: analyses of prognostic factors and recurrence patterns. *Ann Surg Oncol* 2007; 14: 74-83.

Akasu T, Moriya Y, Ohashi Y, Yoshida S, Shirao K, Kodaira S. Adjuvant chemotherapy with uracil-tegafur for pathological stage III rectal cancer after mesorectal excision with selective pelvic lymphadenectomy: a multicentre randomised controlled trial. *Jpn J Clin Oncol* 2006; 36 (4): 237-244.

Akle CA. Endoprotheses for colonic strictures. *Br J Surg* 1998; 85: 310-4.

Allison JE. Review article: faecal occult blood testing for colorectal cancer. *Aliment Pharmacol Ther* 1998; 12: 1-10.

Amado RG, Wolf M, Peeters M, Van CE, Siena S, Freeman DJ et al. Wild-type KRAS is required for panitumumab efficacy in patients with metastatic colorectal cancer. *J Clin Oncol* 2008 April 1; 26 (10): 1626-34.

Amat A, Pescatori M. Perioperative blood transfusions for the recurrence of colorectal cancer. *Cochrane Database Syst Rev* 2006: CD005033.

American Society of Colorectal Surgeons & Wound Ostomy and Continence Nursing Society. ASCRS and WOCN joint position statement on the value of preoperative stoma marking for patients undergoing fecal ostomy surgery. Published 2003. Accessed June 25, 2007.

Andersen HK, Lewis SJ, Thomas S (2006). Early enteral nutrition within 24h of colorectal surgery versus later commencement of feeding for postoperative complications. Cochrane Database Syst RevCD004080.

Andersen J, Hjort-Jakobsen D, Christiansen P, Kehlet H (2007). Readmission rates after planned hospital stay of 2 versus 3 days in fast-track colonic surgery. *Br J Surg* 94: 890-893.

Anderson WF, Guyton KZ, Hiatt RA, Vernon SW, Levin B, Hawk E. Colorectal cancer screening for persons at average risk. *J Natl Cancer Inst* 2002; 94: 1126-33.

Andre T, Boni C, Mounedji-Boudlbf L, Navarro M, Taberbero J, Hickish T et al. "Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer". *N Engl J Med*, Vol. 350, No. 23, 2004, pp. 2343-2351.

Andre T, Quinaux E, Louvet C, Colin P, Gamelin E, Bouche O et al. "Phase III study comparing a semimonthly with a monthly regimen of fluorouracil and leucovorin as adjuvant treatment for stage II and III colon cancer patients: final results of GERCOR C96.1". *J Clin Oncol*, Vol. 25, No. 24, 2007, pp. 3732-3738.

André T, Sargent D, Tarberbero J et al. Current Issues in Adjuvant Treatment of Stage II Colon Cancer. *Ann Surg Oncol* 2006; 13 (6): 887-898.

Anthony T, Fleming JB, Bieligm SC, Sarosi GA, Kim LT, Gregorcyk SG et al. Postoperative colorectal cancer surveillance. *J Am Coll Surg* 2000; 190: 737-749.

Anthony T, Simmang C, Hyman N, Buie D, Kim D, Cataldo P et al. Practice parameters for the surveillance and follow-up of patients with colon and rectal cancer. *Dis Colon Rectum* 2004; 47: 807-817.

Arulampalam T, Costa D, Visvikis D, Boulos P, Taylor I, Ell P. The impact of FDG-PET on the management algorithm for recurrent colorectal cancer. *Eur J Nucl Med* 2001; 28: 1758-1765.

Arumugum PJ, Bevan L, Macdonald L, Watkins AJ, Morgan AR, Beynon J et al. A prospective audit of stomas-analysis of risk factors and complications and their management. *Colorectal Dis* 2003; 5 (1): 49-52.

Arunachalam PS, Putnam G, Jennings P, Messersmith R, Robson AK. Role of computerized tomography (CT) scan of the chest in patients with newly diagnosed head and neck cancers. *Clin Otolaryngol Allied Sci* 2002; 27: 409-411.

Atkin WS, Hart A, Edwards R, Cook CF, Wardle J, McIntyre P et al. Single blind, randomised trial of efficacy and acceptability of oral picolax versus self administered phosphate enema in bowel preparation for flexible sigmoidoscopy screening. *BMJ* 2000; 3: 1504-8.

Atkin W, Rogers P, Cardwell C, Cook C, Cuzick J, Wardle J et al. Wide variation in adenoma detection rates at screening flexible sigmoidoscopy. *Gastroent* 2004; 126: 1247-56.

Augestad KM, Vonen B, Aspevik R, Nestvold T, Ringberg U, Johnson R et al. Should the surgeon or the general practitioner (GP) follow up patients after surgery for colon cancer. A randomized controlled trial protocol focusing on quality of life, cost-effectiveness and serious clinical events. *BMC Health Serv Res* 2008; 8: 137-145.

Barillari P, Ramacciato G, De Angelis R, Gozzo P, Indinnimeo M, Valabrega S et al. Effect of preoperative colonoscopy on the incidence of synchronous and metachronous neoplasms. *Acta Chir Scand* 1990; 156: 163-6.

Barillari P, Ramacciato G, Manetti G, Bovino A, Sammartino P, Stipa V. Surveillance of colorectal cancer: effectiveness of early detection of intraluminal recurrences on prognosis and survival of patients treated for cure. *Dis Colon Rectum* 1996; 39: 388-393.

Barrow E, Alduaij W, Robinson L, Shenton A, Clancy T, Lalloo F et al. Colorectal cancer in HNPCC: Cumulative lifetime incidence, survival and tumour distribution. A report of 121 families with proven mutations. *Clin Genet* 2008.

Bass EM, Del Pino A, Tan A, Pearl RK, Orsay CP, Abcarian H. Does preoperative stoma marking and education by the enterostomal therapist affect outcome? *Dis Colon Rectum* 1997; 40 (4) 400-442.

Basse L, Jakobsen DH, Bardram L, Billesbølle P, Lund C, Mogensen T et al. Functional recovery after open versus laparoscopic colonic resection: a randomized, blinded study. *Ann Surg* 2005; 241: 416-423.

Basse L, Thorbol J, Lossi K, Kehlet H (2004). Colonic surgery with accelerated rehabilitation or conventional care. *Dis Colon Rectum* 47: 271-278.

Basse L, Werner M, Kehlet H (2000). Is urinary drainage necessary during continuous epidural analgesia after colonic resection? *Reg Anesth Pain Med* 25: 498-501. Bastrup 2007.

Beebe DS, McNevin MP, Crain JM, Letourneau JG, Belani KG, Abrams JA et al. Evidence of venous stasis after abdominal insufflation for laparoscopic cholecystectomy. *Surg Gynecol Obstet* 1993; 176: 443-447.

Beets-Tan RG, Beets GL. Rectal cancer: review with emphasis on MR imaging. *Radiology* 2004; 232: 335-46.

Beier-Holgersen R, Boesby S (1996). Influence of postoperative enteral nutrition on postsurgical infections. *Gut* 39; 833-835.

Benoist S, panis Y, Denet C, Mauvais F, Mariani P, Valleur P (1999). Optimal duration of urinary drainage after rectal resection: a randomized controlled trial. *Surgery* 125: 135-141.

Benson AB 3 rd, Schrag D, Somerfield MR et al. American Society of Clinical Oncology recommendations on adjuvant chemotherapy for stage II colon cancer. *J Clin Oncol* 2004; 22: 3408-3419.

Bentrem DJ, Okabe S, Wong WD, Guillem JG, Weiser MR, Temple LK et al. T1 adenocarcinoma of the rectum: transanal excision or radical surgery? *Ann Surg* 2005; 242: 472-7.

Beresford M, Glynne-Jones R et al. The reliability of lymph-node staging in rectal cancer after preoperative chemoradiotherapy. *Clinical Oncology* 2005; 17: 448-55.

Bergmann U, beger HG. Endoscopic mucosal resection for advanced non-polypoid colorectal adenoma and early stage carcinoma. *Surg Endosc* 2003; 17: 475-9.

Bergqvist D, Agnelli G, Cohen AT, Eldor A, Nilsson PE, Moigne-Amrani A et al. Duration of prophylaxis against venous thromboembolism with enoxaparin after surgery for cancer. *N Engl J Med* 2002; 346: 975-980.

Bernstein I and Bisgaard M.L. Årsrapport 2003, HNPCC-registret.

Bernstein I, Bisgaard ML, Myrhøj T. Prevention of colorectal cancer in families with hereditary nonpolyposis colorectal cancer. *Ugeskr Laeger* 2003; 165: 221-225.

Binkert CA, Ledermann H, Jost R, Saurenmann P, Decurtins M, Zollikofer CL. Acute colonic obstruction: clinical aspects and cost-effectiveness of preoperative and palliative treatment with self-expanding metallic stents – a preliminary report. *Radiol* 1998; 206: 199-204.

Bipat S, Glas AS, Slors FJ, Zwinderman AH, Bossuyt PM, Stoker J. Rectal cancer: local staging and assessment of lymph node involvement with endoluminal US, CT, and MR imaging – a meta-analysis. *Radiol* 2004; 232: 773-83.

Bipat S, van Leeuwen MS, Pijl ME, Bossuyt PMM, Zwinderman AH, Stoker J. Colorectal metastases: CT, MR Imaging, and PET for diagnosis – Meta-analysis. *Radiology* 2005; 237: 123-131.

Birgisson H, Talbäck M, Gunnarsson U, Pålman L, Glimelius B. Improved survival in cancer of the colon and rectum in Sweden. *Eur J Surg Oncol* 2005; 31: 845-53.

Bjerkset T, Edna TH. Rectal cancer: the influence of type of operation on local recurrence and survival. *Eur J Surg* 1996; 162: 643-648.

Bjerregaard NC, Tottrup A, Sørensen HT, Laurberg S. Evaluation of the Danish national strategy for selective use of colonoscopy in symptomatic outpatients without known risk factors for colorectal cancer. *Scand J Gastroent* 2007; 42: 228-36.

Blomqvist L, Machado M, Rubio C, Gabrielsson N, Granqvist S, Goldman S et al. Rectal tumour staging: MR imaging using pelvic phased-array and endorectal coils vs endoscopic ultrasonography. *Eur Radiol* 2000; 10: 653-60.

Blyth S, Blakeborough A, Peterson M, Cameron IC, Majeed AW. Sensitivity of magnetic resonance imaging in the detection of colorectal liver metastases. *Ann R Coll Surg Engl* 2008; 90: 25-28.

Bojesen Se, Bernstein I, Gerdes AM, Hartlev M, Koch L, Lindorff-Larsen K et al. Hereditary nonpolyposis colorectal cancer in Denmark – a health technology assessment. Copenhagen: National Board of Health, Danish Centre for health technology assessment 2007; 1-154. Ref Type: Report 2007.

Bonjer HJ, Hop WC, Nelson H, Sargent DJ, Lacy AM, Castelis A et al. Laparoscopically assisted vs open colectomy for colon cancer: a meta-analysis. *Arch Surg* 2007; 142: 298-303.

Borie F, Combescure C, Daures JP, Tretarre B, Millat B. Cost-effectiveness of Two Follow-up Strategies for Curative Resection of Colorectal Cancer: Comparative Study Using a Markov Model. *World J Surg* 2004.

Borowski DW, Kelly SB, Bradburn DM, Wilson RG, Gunn A, Ratcliffe AA et al. Impact of surgeon volume and specialization on short-term outcomes in colorectal cancer surgery. *Br J Surg* 2007; 94: 880-9.

Bosset JF, Calais G, Mineur L, Maingon P, Radosevic-Jelic L, Daban A et al. Enhanced tumoricidal effect of chemotherapy with preoperative radiotherapy for rectal cancer: preliminary results – EORTC 22921. *J Clin Oncol* 2005; 23: 5620-7.

Bosset JF, Collette L, Calais G et al. Chemotherapy with preoperative radiotherapy in rectal cancer. *New Engl j Med* 2006; 355: 1114-1123.

Bowles CJ, Leicester R, Romaya C, Swarbrick E, Williams CB, Epstein O. A prospective study of colonoscopy practice in the UK today: are we adequately prepared for national colorectal cancer screening tomorrow? *Gut* 2004; 53: 277-83.

Bozzetti F, Gavazzi C, Miceli R, Rossi N, Mariani L, Cozzaglio L et al (2000). Perioperative total parenteral nutrition in malnourished, gastrointestinal cancer patients: a randomized, clinical trial. *JPEN J Parenter Enteral Nutr* 24: 7-14.

Brady M, Kinn S, Stuart P (2003). Preoperative fasting for adults to prevent perioperative complications. *Cochrane Database Syst Rev* CD004423.

Braga M, Frasson M, Vignali A, Zuliani W, Capretti G, Di CV. Laparoscopic resection in rectal cancer patients: outcome and cost-benefit analysis. *Dis Colon Rectum* 2007; 50: 464-471.

Braga M, Frasson M, Vignali A, Zuliani W, Civelli V, Di CV. Laparoscopic vs open colectomy in cancer patients: long-term complications, quality of life, and survival. *Dis Colon Rectum* 2005; 48: 2217-2223.

Branagan G, Finnis D, Wessex Colorectal Cancer Audit Working Group. Prognosis after anastomotic leakage in colorectal surgery. *Dis Colon Rectum* 2005 May; 48 (5): 1021-6.

Brandstrup B, Tonnesen H, Beier-Holgersen R, Hjortso E, Ording H, Lindorff-Larsen K et al (2003). Effects of intravenous fluid restriction on postoperative complications: comparison of two perioperative fluid regimens: a randomized assessor-blinded multicenter trial. *Ann Surg* 238: 641-648.

Bretagnol F, Alves A, Ricci A, Valleur P, Panis Y (2007). Rectal cancer surgery without mechanical bowel preparation. *Br J Surg* 1994: 1266-1271.

Bretagnol F, Rullier E, Laurent C, Zerbib F, Gontier R, Saric J. Comparison of functional results and quality of life between intersphincteric resection and conventional coloanal anastomosis for low rectal cancer. *Dis Colon Rectum* 2004; 47: 832-8.

Bretagnol F, Troubat H, Laurent C, Zerbib F, Saric JU, Rullier E. Long-term functional results after sphincter-saving resection for rectal cancer. *Gastroenterol Clin Biol* 2004; 28: 155-9.

Bretthauer M, Thiis-Evensen E, Huppertz-Hauss G, Gisselsson L, Grotmol T, Skovlund E et al. NORCCAP (Norwegian colorectal cancer prevention): a randomised trial to assess the safety and efficacy of carbon dioxide versus air insufflation in colonoscopy. *Gut* 2002; 50: 604-7.

Breukink S, Pierie J, Wiggers T. Laparoscopic versus open total mesorectal excision for rectal cancer. *Cochrane Database Syst Rev* 2006 (4): CD005200.

Brown G, Richards CJ, Newcombe RG, Dallimore NS, Radcliffe AG, Carey DP. Rectal carcinoma: thin-section MR imaging for staging in 28 patients. *Radiol* 1999; 211: 215-22.

Brown G, Kirkham A, Williams GT, Bourne M, Radcliffe AG, Sayman J et al. High-resolution MRI of the anatomy important in total mesorectal excision of the rectum. *Am J Roentgenol* 2004; 182: 431-9.

Brown G, Daniels IR, Richardson C, Revell P, Peppercorn D, Bourne M. Techniques and trouble-shooting in high spatial resolution thin slice MRI for rectal cancer. *Br J Radiol* 2005; 78: 245-51.

Bruinvels DJ, Stiggelbout AM, Kievit J, van Houwelingen HC, Habbema JD, van de Velde CJ. Follow-up of patients with colorectal cancer. A Meta-analysis. *Ann Surg* 1994; 219: 174-182.

Bucher P, Gervaz P, Soravia C, Mermillod B, Erne M, Morel P (2005). Randomized clinical trial of mechanical bowel preparation versus no preparation before elective left-sided colorectal surgery. *Br J Surg* 92: 409-414.

Bucher P, Mermillod B, Gervaz P, Morel P (2004). Mechanical bowel preparation for elective colorectal surgery: a meta-analysis. *Arch Surg* 139: 1359-1364.

Buess G, Hutterer F, Theiss J, Böbel M, Isselhard W, Pichlmaier H. A system for a transanal endoscopic rectum operation. *Chir* 1984; 55: 677-80.

Buhre LM, Plukker JT, Mehta DM, Verschueren RC, Oldhoff J. The extended Hartmann operation as an elective procedure for rectal cancer. A forgotten operation. *Eur J Surg Oncol* 1991; 17: 502-6.

Bujko K, Nowacki MP, Nasierowska-Guttmejer A, Michalski W, Bebenek M, Kryj M. Long-term results of a randomized trial comparing preoperative short-course radiotherapy with preoperative conventionally fractionated chemoradiation for rectal cancer. *Br J Surg* 2006; 93: 1215-23.

Bujko K, Kepka L, Michalski W, Nowacki M. Does rectal cancer shrinkage induced by preoperative radio(chemo)therapy increase the likelihood of anterior resection? A systematic review of randomised trials. *Radiother Oncol* 2006; 80 (1): 4-12.

Bundgaard-Nielsen M, Holte K, Secher N, Kehlet H (2007). Monitoring of peri-operative fluid administration by individualized goal-directed therapy. *Acta Anaesthesiol Scand* 51: 331-340.

Burke P, Mealy K, Gillen P, Joyce W, Traynor O, Hyland J (1994). Requirement for bowel preparation in colorectal surgery. *Br J Surg* 81: 907-910.

Burt RW. Colon cancer screening. *Gastroenterology* 2000; 119: 837-853.

Burton S, Brown G, Daniels IR. MRI directed multidisciplinary team preoperative strategy the way to eliminate positive circumferential margins? *Br J Cancer* 2006; 94: 351-7.

Bussey HJ, Wallace MH, Morson BC. Metachronous carcinoma of the large intestine and intestinal polyps. *Proc R Soc Med* 1967; 60: 208-210.

Bülow S, Christensen IJ. Sigmoidoskopi som primær undersøgelse ved tarmsymptomer. *Ugeskr Laeger* 2001; 19: 6573-6.

Bülow S, Björk J, Christensen IJ, Fausa O, Järvinen H, Moesgaard F. Duodenal adenomatosis in familial adenomatous polyposis. *Gut* 2004; 53: 381-6.

Bülow S, Bülow C, Björk J, Järvinen H, Vasen HFA, Christensen IJ. Colectomy and ileorectal anastomosis is still an option in selected patients with familial adenomatous Polyposis. *Dis Colon Rectum* 2008; 51: 1318-23.

Bülow S, Jensen LH, Altaf R, Harling H, Jensen M et al. A national cohort study of long course preoperative radiotherapy in primary fixed rectal cancer in Denmark. *Colorectal Dis* 2009, I trykken.

Bülow S, Moesgaard FA, Billesbølle P et al. Anastomoselækage efter lav anterior resektion for rectumcancer. *ugeskr Læger* 1997; 159: 297-301.

Bülow S. Familiær adenomatøs polypose. FAP. Vejledende retningslinjer for diagnose, behandling og control. Polyposeregistret, Hvidovre Hospital & Danish Colorectal Cancer Group. 2. udgave. Hvidovre Hospital, 2003.

Bülow S. FAP. Familiær adenomatøs polypose. Information for polyposepatienter og deres familie. Eds: Holten I, Munk TB, Olsen T. 3. udgave. *Kræftens Bekæmpelse* 2007 og www.cancer.dk.

Böhm B, Schwenk W, Huckle HP, Stock W. Does methodic long-term follow-up affect survival after curative resection of colorectal carcinoma? *Dis Colon Rectum* 1993; 36: 280-286.

Cahill RA. What's wrong with sentinel node mapping in colonic cancer. *World J Gastroenterology* 2007; 13 (47): 6291-94.

Cali RL, Pitsch RM, Thorson AG, Watson P, Tapia P, Blatchford GJ et al. Cumulative incidence of metachronous colorectal cancer. *Dis Colon Rectum* 1993; 36: 388-393.

Camilleri-Brennan J, Steele RJ. Quality of life after treatment for rectal cancer. *Br J Surg* 1998; 85: 1036-43.

Camilleri-Brennan J, Steele RJ. Objective assessment of morbidity and quality of life after surgery for low rectal cancer. *Colorectal Dis* 2002; 4: 61-66.

Capirci C, Valentini V, Rodel F et al. Prognostic value of pathological complete response after neoadjuvant radiochemotherapy in locally advanced rectal cancer: a pooled analysis on 566 patients. *Radiother Oncol* 2006; 81 (Suppl 1): S76-S77 (abstracts 193).

Carmody BJ, Otchy DP. Learning curve of transrectal ultrasound. *Dis Colon Rectum* 2000; 43: 193-7.

Cassidy J, Clarke S, az-Rubio E, Scheithauer W, Figer A, Wong R et al. Randomized phase III study of capecitabine plus oxaliplatin compared with fluorouracil (folinic acid plus oxaliplatin as first-line therapy for metastatic colorectal cancer. *J Clin Oncol* 2008 April 20; 26 (12): 2006-12.

Cecil TD, Taffinder N, Gudgeon AM. A personal view on laparoscopic rectal cancer surgery. *Colorectal Dis* 2006; 8 Suppl 3: 30-32.

Chan JA, Meyerharadt JA, Niedzwiecki D, Hollis D, Saltz LB, Mayer RJ et al. Association of family history with cancer recurrence and survival among patients with stage III colon cancer. *Jama* 2008; 299: 2515-2523.

Chaudhri S, Brown L, Hassan I, Horgan AF. Preoperative intensive, community based vs traditional stoma education: a randomized, trolled trial. *Dis Colon Rectum* 2005; 48 (3): 504-509.

Chiu HM, Chang CY, Chen CC, Lee YC, Wu MS, Lin JT et al. Preparation for colonoscopy in hospitalized patients. *Dig Dis Sci* 2007; 52: 835-9.

Cho YB, Lee WY, Yun HR, Lee WS, Yun SH, Chun HK. Tumor localization for laparoscopic colorectal surgery. *World J Surg* 2007; 31: 1491-1495.

Choi YJ, Park SH, Lee SS, Choi EK, Yu CS, Kim HC et al. CT colonography for follow-up after surgery for colorectal cancer. *Am J Roentgenol* 2007; 189: 283-289.

Chorev N, Chadad B, Segal N, Shemesh I, Mor M, Plaut S et al. Preparatoin for colonoscopy in hospitalized patients. *Dig Dis Sci* 2007; 52: 835-9.

Choy PYG, Bisset IP, Docherty JG, Parry BR, Merrie AEH. Stapled versus handsewn methods for ileocolic anastomoses. *Cochrane Database of Systematic Reviews* 2007, Issue 3. Art.No.: CD004320.

Church JM. Analysis of the colonoscopic findings in patients with rectal bleeding according to the pattern of their presenting symptoms. *Dis Colon Rectum* 1991; 34: 391-5.

Church JM. Experience in the endoscopic management of large polyps. *ANZ J Surg* 2003 Dec; 73 (12): 988-95.

Collette L, Bosset JF, den Dulk M et al. Patients With Curative Resection of cT3-4 Rectal Cancer After Preoperative Radiotherapy or Radiochemo: Does Anybody Benefit From Adjuvant Fluorouracil-Based Chemotherapy? A Trial of the European Organisation for Research and Treatment of Cancer Radiation Oncology Group. *J Clin Oncol* 2007; 25: 4379-4386.

Collins R, Scrimgeour A, Yusuf S, Peto R. Reduction in fatal pulmonary embolism and venous thrombosis by perioperative administration of subcutaneous heparin. Overview of results of randomized trials in general, orthopaedic, and urologic surgery. *N Engl J Med* 1988; 318: 1162-1173.

Colorectal Cancer Collaborataive Group. Adjuvant radiotherapy for rectal cancer: a systematic overview of 22 randomized trials involving 8507 patients. *Lancet* 2001; 358: 1291-1304.

Colorectal Cancer Collaborative Group. Palliative chemotherapy for advanced or metastatic colorectal cancer. Colorectal Meta-analysis Collaboration. Cochrane Database Syst Rev 2000; (2): CD001545.

Comella P, Massidda B, Filippelli G, Palmeri S, Natale D, Farris A et al. Oxaliplatin plus high-dose folinic acid and 5-fluorouracil iv bolus (OXAFUFU) versus irinotecan plus high-dose folinic acid and 5-fluorouracil iv bolus (IRIFUFU) in patients with metastatic colorectal carcinoma: a Southern Italy Cooperative Oncology Group phase III trial. Ann Oncol 2005 June; 16 (6): 878-86.

Compton C, Fenoglio-Preiser CM, Peeigrew N et al. American Joint Committee on Cancer Prognostic Factors Consensus Conference: Colorectal Working Group. Cancer 2000; 88: 1739-1757.

Contant CM, Hop WC, van't Sant HP, Oostvogel HJ, Smeets HJ, Stassen LP et al (2007). Mechanical bowel preparation for elective colorectal surgery: a multicentre randomised trial. Lancet 370: 2112-2117.

Cook AD, Single R, McCahill LE. Surgical Resection of Primary Tumors in Patients Who Present With Stage IV Colorectal Cancer: An Analysis a Surveillance, Epidemiology, and End Results Data, 1988 to 2000. Ann Surg Oncol 2005; 12 (8): 637-645.

Cornish JA, Tilney HS, Heriot AG, Lavery IC, Fazio VW, Tekkis PP. A meta-analysis of quality of life for abdominoperineal excision of rectum versus anterior resection for rectal cancer. Ann Surg Oncol 2007; 14: 2056-68.

Crowe PJ, Temple WJ, Lopez MJ, Ketchamas AS. Pelvic Excenteration for Advanced Pelvic Malignancy. Sem Surg Oncol 1999; 17: 152-60.

Cunningham D, Humblet Y, Siena S, Khayat D, Bleiberg H, Santoro A et al. Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer. N Engl J Med 2004 July 22; 351 (4): 337-45.

Curti G, Maurer CA, Büchler MW. Colorectal carcinoma: is lymphadenectomy useful? Dig Surg 1998; 15: 193-208.

Dahlberg M et al. Preoperative irradiation affects functional results after surgery for rectal cancer: results from a randomized study. Dis Colon Rectum 1998 May; 41 (5): 543-9; discussion 549-51.

Daniels IR, Woodward S, Taylor FG, Raja A, Toomey P. Female urogenital dysfunction following total mesorectal excision for rectal cancer. World J Surg Oncol 2006; 4: 6.

de Gramont A, Bosset JF, Milan C, Rougier P, Bouche O, Etienne PL et al. Randomized trial comparing monthly low-dose leucovorin and fluorouracil bolus with bimonthly high-dose leucovorin and fluorouracil bolus plus continuous infusion for advanced colorectal cancer: a French intergroup study. J Clin Oncol 1997 February; 15 (2): 808-15.

de Gramont A, Figer A, Seymour M, Homerin M, Hmissi A, Cassidy J et al. Leucovorin and fluorouracil with or without oxaliplatin as first-line treatment in advanced colorectal cancer. *J Clin Oncol* 2000 August; 18 (16): 2938-47.

De Salvo GL, Gava C, Lise M et al. Curative surgery for obstruction from primary left colorectal carcinoma: Primary or staged resection. *The Cochrane Database of Systemic Reviews* 2006 Issue 2: The Cochrane Collaboration, John Wiley et Sons, Ltd.

de Silva WM. Developing a molecular marker for metachronous colorectal cancer. *Ceylon Med J* 1999; 44: 162-165.

de Vos tot Nederveen Cappel WH, Buskens E, van Duijvendijk P, Menko FH, Griffioen G, Slors JF et al. Decision analysis in the surgical treatment of colorectal cancer due to mismatch repair gene defect. *Gut* 2003; 52 (12): 1752-55.

de Vos tot Nederveen Cappel WH, Nagengast FM, Griffioen G, Menko FH, Taal BG, Kleiburker JH et al. Surveillance for hereditary nonpolyposis colorectal cancer: A long term study on 114 families. *Dis Colon Rectum* 2002; 45 (12): 1588-94.

den Dulk M, Marijnen CA, Putter H, Rutten HJ, Beets GL, Wiggers T et al. Risk factors for adverse outcome in patients with rectal cancer treated with an abdominoperineal resection in the total mesorectal excision trial. *Ann Surg* 2007; 246: 83-90.

den Dulk M, Marijnen CAM, Putter H. Risk factors for adverse outcome in patients with rectal cancer treated with an abdominoperineal resection in the Total Mesorectal Excision trial. *Ann Surg* 2007; 246: 83-90.

Desch CE, Benson AB 3rd, Smith TJ, Flynn PJ, Krause C, Loprinzi CL. Recommended colorectal cancer surveillance guidelines by the American Society of Clinical Oncology. *Clin Oncol* 1999; 17: 1312.

Desch CE, Benson AB III, Somerfield MR, Flynn PJ, Krause C, Loprinzi CL et al. Colorectal cancer surveillance: 2005 update of an American Society of Clinical Oncology Practice guideline. *J Clin Oncol* 2005; 23: 8512-8515.

Desnoo L, Faithfull S. A qualitative study of anterior resection syndrome: the experiences of cancer survivors who have undergone resection surgery. *Eur J Cancer Care* 2006; 15: 244-51.

Detsky AS, Baker Jp, O'Rourke K, Johnston N, Whitwell J, Mendelson RA et al (1987). Predicting nutrition-associated complications for patients undergoing gastrointestinal surgery. *JPEN J Parenter Enteral Nutr* 11: 440-446.

Di Fiore F, Van Cutsem E, Laurent-Puig P, Siena S, Frantini M, De Roock W et al. Role of KRAS mutation in predicting response, progression-free survival, and overall survival in irinotecan-refractory patients treated with cetuximab plus irinotecan for a metastatic colorectal cancer: Analysis of 281 individual data from published series. *Proc Am Soc Clin Oncol* 2008; #4035.

Dietrich CF, Kratzer W, Strobe D et al. Assessment of metastatic liver disease in patients with primary extrahepatic tumours by contrast-enhanced sonography versus CT and MRI. *World J Gastroenterol* 2006; 12: 1699-1705.

Dincler S, Koller MT, Steurer J, Bachmann LM, Christen D, Buchmann P. Multidimensional analysis of learning curves in laparoscopic sigmoid resection: eight-year results. *Dis Colon Rectum* 2003; 46: 1371-1378.

Doria-Rose VP, Newcomb PA, Levin TR. Incomplete screening flexible sigmoidoscopy associated with female sex, age, and increased risk of colorectal cancer. *Gut* 2005; 54: 1273-8.

Dostalík J, Martinek L, Vavra P, Andel P, Gunka I, Gunkova P. Laparoscopic colorectal surgery in obese patients. *Obes Surg* 2005; 15: 1328-1331.

Douillard JY, Cunningham D, Roth AD, Navarro M, James RD, Karasek P et al. Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: a multicentre randomised trial. *Lancet* 2000 March 25; 355 (9209): 1041-7.

Douillard JY, Hoff PM, Skillings JR, Eisenberg P, Davidson N, Harper P et al. Multicenter phase III study of uracil/tegafur and oral leucovorin versus fluorouracil and leucovorin in patients with previously untreated metastatic colorectal cancer. *J Clin Oncol* 2002 September 1; 20 (17): 3605-16.

Dove-Edwin I, Sasieni P, Adams J, Thomas HJ. Prevention of colorectal cancer by colonoscopic surveillance in individuals with family history of colorectal cancer: 16 year, prospective, follow-up study. *BMJ* 2005; 331: 1047-1052.

Dubé S, Heyen F, Jenicek M. Adjuvant chemotherapy in colorectal carcinoma: results of a meta-analysis. *Dis Colon Rectum* 1997; 40 (1): 35-41.

Dunlop MG. Guidance Dunlop MG. Guidance on large bowel surveillance for people with two first degree relatives with colorectal cancer or one first degree relative diagnosed with colorectal cancer under 45 years. *Gut* 2002; 51 suppl 5: V17-V20.

Emborg H, Heuer O, Larsen P. 2006. DANMAP 2004. DTU Fødevareinstituttet, DTU Veterinærinstituttet.

Engel C, Forberg J, Holinski-Feder E, Pagenstecher C, Plaschke J, Kloor M et al. Novel strategy for optimal sequential application of clinical criteria, immunohistochemistry and microsatellite analysis in the diagnosis of hereditary nonpolyposis colorectal cancer. *Int J Cancer* 2006; 118: 115-122.

Engelen SM, Beets GL, Beets-Tan RG. Role of preoperative local and distant staging in rectal cancer. *Onkologie* 2007; 30: 141-5.

Enoxacan Study Group. Efficacy and safety of enoxaparin versus unfractionated heparin for prevention of deep vein thrombosis in elective cancer surgery: A double-blind randomised multicentre trial with venographic assessment. *Br J Surg* 1997; 84: 1099-1103.

Ewald J, Rodrigue CM, Mourra N, Lefevre JH, Flejou JF, Tiret E et al. Immunohistochemical staining for mismatch repair proteins, and its relevance in the diagnosis of hereditary non-polyposis colorectal cancer. *Br J Surg* 2007; 94: 1020-1027.

Faerden AE, Naimy N, Wiik P, Reiertsen O, Weyessa S, Trønnes S et al. Total mesorectal excision for rectal cancer: difference in outcome for low and high rectal cancer. *Dis Colon Rectum* 2005; 48: 2224-31.

Fa-Si Oen P, Roumen R, Buitenweg J, van DV, van GD, Putter H et al (2005). Mechanical bowel preparation or not? Outcome of a multicenter, randomized trial in elective open colon surgery. *Dis Colon Rectum* 48: 1509-1516.

Farouk R, Ratnaval CD, Monson JRT, Lee PWR. Staged delivery of Nd: YAG laser therapy for palliation of advanced rectal carcinoma. *Dis Colon Rectum* 1997; 40: 156-60.

Fazio VW. Indications and surgical alternatives for palliation of rectal cancer. *J Gastrointest Surg* 2004; 8: 262-5.

Fazio VW, Zutshi M, Remzi FH, Parc Y, Ruppert R, Fürst A et al. A randomized multicenter trial to compare long-term functional outcome, quality of life, and complications of surgical procedures for low rectal cancers. *Ann Surg* 2007; 246:L 481-8.

Felsher JJ, Olesevich M, Farres H, Rosen M, Fanning A, Dunkin BJ et al. Validation of a flexible endoscopy simulator. *Am J Surg* 2005; 189: 487-500.

Feng YS, Hsu H, Chen SS. One-stage operation for obstructing carcinomas of the left colon and rectum. *Dis Colon Rectum* 1987; 30: 29-32.

Ferenschild FTJ, Vermaas M, Nuyttens JJME, Graveland WJ, Marinelli AWKS, van der Sijp JR et al. Value of Intraoperative Radiotherapy in Locally Advanced Rectal Cancer. *Dis Colon Rectum* 2006; 49: 1257-65.

Fernandez-Represa JA, Mayol JM, Garcia-Aguilar J. Total mesorectal excision for rectum cancer: the truth lies underneath. *World J Surg* 2004; 28: 113-6.

Fielding LP, Stewart-Brown S, Blesovsky L. Large-bowel obstruction caused by cancer: a prospective study. *Br Med J* 1979; 2: 515-7.

Fielding LP, Stewart-Brown S, Blesovsky L, Kearney G. Anastomotic integrity after operations for large-bowel cancer: a multicentre study. *Br Med J* 1980; 281: 411-4.

Fielding LP, Phillips RK, Fry JS, Hittinger R. Prediction of outcome after curative resection for large bowel cancer. *Lancet* 1986; 2: 904-7.

Fielding LP, Phillips RK, Hittinger R. Factors influencing mortality after curative resection for large bowel cancer in elderly patients. *Lancet* 1989; 1: 595-7.

Fietkau R, Barten M, Klautke G et al. Postoperative chemotherapy may not be necessary for patients with ypN0-category after neoadjuvant chemoradiotherapy of rectal cancer. *Dis Colon Rectum* 2006; 49 (9): 1284-1292.

Figueredol A, Rumble RB, Maroun J, Earle CC, Cummings B, McLeod R et al. Wollow-up of patients with curatively resected colorectal cancer: a practice guideline. *BMC Cancer* 2003; 3: 26.

Fisher B, Wolmark N, Rockette H et al. Postoperative adjuvant chemotherapy or radiation therapy for rectal cancer: results from NSABP protocol R-01. *J Natl Cancer Inst* 1988; 80 (1): 21-9.

Fleshman J, Sargent DJ, Green E, Anvari M, Stryker SJ, Beart RW Jr et al. Laparoscopic colectomy for cancer is not inferior to open surgery based on 5-year data from COST Study Group trial. *Ann Surg* 2007; 246: 655-662.

Folkersen S, Jakobsen DH, Andreassen S, Basse L, Kehlet H (2005). Accelererede colon kirurgiske forløb. MTV-rapport. Sundhedsstyrelsen.

Folprecht G, Seymour MT, Saltz L, Douillard JY, Stephens RJ, Van Cutsem E et al. Irinotecan/5-FU/FA (I-FU) or 5-FU/FA (FU) first-line therapy in older and younger patients with metastatic colorectal cancer: Combined analysis of 2,691 patients in randomized controlled trials. *Proc Am Soc Clin Oncol* 2007; #4071.

Fong Y et al. Clinical Score for Predicting Recurrence after Hepatic Resection for Metastatic Colorectal Cancer. Analysis of 1001 Consecutive Cases. *Ann Surg* 1999; 230: 309-21.

Fong Y, Saldinger PF, Akhurst T et al. Utility of 18F-FDG positron emission tomography scanning on selection of patients for resection of hepatic colorectal metastases. *Am J Surg* 1999; 178: 282-287.

Francini G, Petrioli R, Lorenzini L, Mancini S, Armenio S, Tanzini G et al. "Folinic acid and 5-fluorouracil as adjuvant chemotherapy in colon cancer". *Gastroenterology*, Vol. 106, No. 4, 1994, pp. 899-906.

Fremstidig organisering af kirurgien – faglighed og sammenhæng. Amdsrådsforeningen, 2005.

Frimodt-Møller N 2002. How predictive is PK/PD for antibacterial agents? *Int J Antimicrob Agents* 19: 333-339.

Fukami N, Lee JH. Endoscopic treatment of large sessile and flat colorectal lesions. *Curr Opin Gastroenterol* 2006 Jan; 22 (1): 54-9.

Galandiuk S, Wieand HS, Moertel CG, Cha SS, Fitzgibbons RJ, Pemberton JH et al. Patterns of recurrence af curative resection of carcinoma of the colon and rectum. *Surg Gyn & Obstetrics* 1992; 174: 27-32.

Gamagami RA, Liagre A, Chiotasso P, Istvan G, Lazorthes F. Goloanal anastomosis for distal third rectal cancer: prospective study of oncologic results. *Dis Colon Rectum* 1999; 42: 1272-5.

Gamagami R, Istvan G, Cabarrot P, Liagre A, Chiotasso P, Lazorthes F. Fecal continence following partial resection of the anal canal in distal rectal cancer: long-term results after coloanal anastomoses. *Surg* 2000; 127: 291-5.

Gan S, Wilson K, Hollington P. Surveillance of patients following surgery with curative intent for colorectal cancer. *World J Gastroenterol* 2007; 13: 3816-3823.

Garcea G et al. Review. Inoperable colorectal liver metastases: A declining entity? *Eur J Cancer* 2008; 44: 2555-72.

Garcia-Osogobio SM, Takahashi Monroy T, Velasco L, Gaxiola M, Sotres Vega A, Santillán-Doherty P. *Rev Invest Clin* 2006 May-Jun; 58 (3): 198-203.

Gastinger I, Marusch F, Steinert R, Wolff S, Koeckerling F, Lippert H et al. Protective defunctioning stoma in low anterior resection for rectal carcinoma. *Br J Surg* 2005; 92: 1137-42.

Gavagan JA, Whiteford MH, Swanstrom LL. Full-thickness intraperitoneal excision by transanal endoscopic microsurgery does not increase short-term complications. *Am J Surg* 2004; 187: 630-4.

Geerts WH, Pineo GF, Heit JA, Bergqvist D, Lassen MR, Colwell CW, Ray JG. Prevention of venous thromboembolism: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004; 126: 338S-400S.

Gennemførlighedsundersøgelse af screening for tarmkræft I Københavns amt 2007.

Gerard JP, Conroy T, Bonnetain F, Bouché O, Chapet O, Closon-Dejardin MT et al. Preoperative radiotherapy with or without concurrent fluororacil and leucovorin in T3-4 rectal cancers: results of FFCD 9203. *J Clin Oncol* 2006; 24: 4620-5.

Gertsch P, Baer HU, Kraft R, Maddern GJ, Altermatt HJ. Malignant cells are collected on circular staplers. *Dis Colon Rectum* 1992; 35: 238-41.

Giacchetti S, Perpoint B, Zidani R, Le Bail N, Faggiuolo R, Focan C et al. Phase III multicenter randomized trial of oxaliplatin added to chronomodulated fluorouracil-leucovorin as first-line treatment of metastatic colorectal cancer. *J Clin Oncol* 2000 January; 18 (1): 136-47.

Giantonio BJ, Catalano PJ, Meropol NJ, O'Dwyer PJ, Mitchell EP, Alberts SR et al. Bevacizumab in combination with oxaliplatin, fluorouracil, and leucovorin (FOLFOX4) for previously treated metastatic colorectal cancer: results from the Eastern Cooperative Oncology Group Study E3200. *Clin Oncol* 2007; 25: 1539-44.

Gill S, Loprinzi CL, Sargent DJ et al. Pooled analysis of fluorouracil-based adjuvant therapy for stage II and III colon cancer: Who benefits and by how much? *J Clin Oncol* 2004; 22: 1797-1806.

Ginnerup Pedersen B, Rosenkilde M, Christiansen TE, Laurberg S. Extracolonic findings at computed tomography colonography are a challenge. *Cut* 2003; 52: 1744-7.

Glimelius B, Jakobsen A, Graf W, Berglund A, Gadeberg C, Hansen P et al. Bolus injection (2-4 min) versus short-term (10-20 min) infusion of 5-fluorouracil in patients with advanced colorectal cancer: a prospective randomised trial. Nordic Gastrointestinal Tumour Adjuvant Therapy Group. *Eur J Cancer* 1998 April; 34 (5): 674-8.

Glimelius B. A systematic review of radiation therapy effects in rectal cancer. *Acta Oncol* 2003; 42: 476-92.

Glover C, Douse P, Kane P et al. Accuracy of investigations for asymptomatic colorectal liver metastases. *Dis Colon Rectum* 2002; 45: 476-484.

Goldberg RM, Fleming TR, Tangen CM, Moertel CG, Macdonald JS, Haller DG et al. Surgery for recurrent colon cancer: strategies for identifying resectable recurrence and success rates after resection. Eastern Cooperative Oncology Group, the North Central Cancer Treatment Group, and the Southwest Oncology Group. *Ann Intern Med* 1998; 129: 27-35.

Goldberg RM, Koehne CH, Seymour MT, de Gramont A, Porschen R, Saltz L et al. A pooled safety and efficacy analysis examining the effect of performance status (PS) on outcomes in nine first-line treatment (rx) trials (cts) of 6,286 patients (pts) with metastatic colorectal cancer (MCR). *Proc Am Soc Clin Oncol* 2007; #4011.

Goldberg RM, Sargent DJ, Morton RH, Fuchs CS, Ramanathan RK, Williamson SK et al. A randomized controlled trial of fluorouracil plus leucovorin, irinotecan, and oxaliplatin combinations in patients with previously untreated metastatic colorectal cancer. *J Clin Oncol* 2004 January 1; 22 (1): 23-30.

Gould MK, Maclean CC, Kuschner WG, Rydzak CE, Ovens DK. Accuracy of positron emission tomography for diagnosis of pulmonary nodules and mass lesions: a metaanalysis. *JAMA* 2001; 285: 914-924.

Gray R, McConkey C. Authors reply. *Lancet* 2008; 371: 1503.

Grossmann EM, Johnson FE, Virgo KS, Longo WE, Fossati R. Follow-up of the colorectal cancer patients after resection with curative intent: the GILDA trial. *Surg Oncol* 2004; 13: 119-124.

Grothey A, Sargent D, Goldberg RM, Schmoll HJ. Survival of patients with advanced colorectal cancer improves with the availability of fluorouracil-leucovorin, irinotecan, and oxaliplatin in the course of treatment. *J Clin Oncol* 2004 April 1; 22 (7): 1209-14.

Grunfeld E, Mant D, Yudkin P, Adewuyi-Dalton R, Cole D, Stewart J et al. Routine follow-up of breast cancer in primary care. randomised trial. *BMJ* 1996; 313: 665-669.

Gualdi GF, Casciani E, Guadataro A, d'Orta C, Polettini E, Pappatardo G. Local staging of rectal cancer with transrectal ultrasound and endorectal magnetic resonance imaging: comparison with histologic findings. *Dis Colon Rectum* 2000; 43: 338-45.

Guenaga KF, Matos D, Castro AA, Atallah AN, Wille-Jørgensen P (2003). Mechanical bowel preparation for elective colorectal surgery. *Cochrane Database Syst Rev* CK001544. Guidelines 2001.

Guillem JF, Chessin DB, Cohen AM et al. Long term oncologic outcome following preoperative combined modality therapy and total mesorectal excision of locally advanced rectal cancer. *Ann Surg* 2005; 241: 829-836.

Guillem JG, Chessin DB, Shia J, Suriawinata A, Riedel E, Moore HG et al. A prospective pathologic analysis using whole-mount sections of rectal cancer following preoperative combined modality therapy: implications for sphincter preservation. *Ann Surg* 2007; 245: 88-93.

Guillou PJ, Quirke P, Thorpe H, Walker J, Jayne DG, Smith AM et al. Short-term endpoints of conventional versus laparoscopic-assisted surgery in patients with colorectal cancer (MRC CLASICC trial): multicentre, randomised controlled trial. *Lancet* 2005; 365: 1718-1726.

Guittet L, Bouvier V, Mariotte N, Vallee JP, Arsene D, Boutreux S et al. Comparison of a guaiac based and an immunochemical faecal occult blood test in screening for colorectal cancer in a general average risk population. *Gut* 2007; 56: 210-4.

Gulbiniene J, Markelis R, Tamelis A, Saladzinskas Z. The impact of preoperative stoma siting and stoma care education on patient's quality of life. *Medicina (Kaunas, Lithuania)* 2004; 40 (11): 1045-1053.

Haider MA, Amitai MM, Rappaport DC et al. Multi-detector row helical CT in preoperative assessment of small (< or =1.5 cm) liver metastases: is thinner collimation better? *Radiology* 2002; 225: 137-142.

Halefoglu AM, Yildirim S, Avlanmis O, Sakiz D, Baykan A. Endorectal ultrasonography versus phase-array magnetic resonance imaging for preoperative staging of rectal cancer. *World J Gastroenterol* 2008; 14: 3504-10.

Halvorsen TB, Seim E. Degree of differentiation in colorectal adenocarcinomas: A multivariate analysis of the influence of survival. *J Clin Pathol* 1988; 41: 532-7.

Hampel H, Frankel WL, Martin E, Arnold M, Khanduja K, Kuebler P et al. Screening for the Lynch syndrome (hereditary nonpolyposis colorectal cancer). *N Engl J Med* 2005; 352 (18): 1851-1860.

Hanson JM, Atkin WS, Cunliffe WJ, Browell DA, Griffith CD, Varma JS et al. Rectal retroflexion: an essential part of lower gastrointestinal endoscopic examination. *Dis Colon Rectum* 2001; 44: 1706-8.

Hara AK, Johnson CD, MacCarty RL, Welch TJ. Incidental extracolonic findings at CT colonography. *Radiol* 2000; 215: 353-7.

Hardcastle JD, Chamberlain JO, Robinson MH, Moss SM, Amar SS, Balfour TW et al. Randomised controlled trial of faecal-occult-blood screening for colorectal cancer. *Lancet* 1996; 348: 1472-7.

Hardcastle JD, Thomas WM, Chamberlain J, Pye G, Sheffield J, James PH et al. Results for first 107,349 subjects. *Lancet* 1989; 1: 1160-4.

Hares MM, Alexander-Williams J (1982). The effect of bowel preparation on colonic surgery. *World J Surg* 6: 175-181.

Harling H, Bülow S, Kronborg O et al. Survival of rectal cancer patients in Denmark during 1994-99. *Colorectal Dis* 2004; 6: 153-7.

Hassan C, Zullo MD, Risio M, Rossini FP, Morini S. Histological risk factors and clinical outcome in colorectal malignant polyp: A pooled-data analysis. *Dis Colon Rectum* 2005; 48: 1588-96.

Hassan I, Larson DW, Cima RR, Gaw JU, Chus HK, Hahnloser D et al. Long-term functional and quality of life outcomes after coloanal anastomosis for distal rectal cancer. *Dis Colon Rectum* 2006; 49: 1266-74.

Haugen V, Bliss DZ, Savik K. perioperative factors that affect long-term adjustment to an ostomy. *J Wound Ostomy Continence Nurs* 2006; 33 (5): 525-535.

Hausel J, Nygren J, Lagerkranser M, Hellstrom PM, Hammarqvist F, Almstrom C et al (2001). A carbohydrate-rich drink reduces preoperative discomfort in elective surgery patients. *Anesth Analg* 93: 1344-1350.

Heah SM, Seow-Choen F, Eu KW, Ho YH, Tang CL. Prospective, randomized trial comparing sigmoid vs. descending colonic J-pouch after total rectal excision. *Dis Colon Rectum* 2002; 45: 322-8.

Heald RJ, Husband EM, Ryall RD. The mesorectum in rectal cancer surgery – the clue to pelvic recurrence? *Br J Surg* 1982; 69: 613-6.

Heald RJ, Ryall RD. Recurrence and survival after total mesorectal excision for rectal cancer. *Lancet* 1986; 1: 1479-82.

Heald RJ, Moran BJ, Ryall RDH, Sexton R, MacFarlane JK. Rectal cancer. The Basingstoke experience of total mesorectal excision, 1978-1997. *Arch Surg* 1998; 133: 894-9.

Heald RJ, Karanjia ND. Results of radical surgery for rectal cancer. *World J Surg* 1992; 16: 848-57.

Heald RJ, Smedh RK, Kald A, Sexton R, Moran BJ. Abdominoperineal excision of the rectum-an endangered operation. Norman Nigro Lectureship. *Dis Colon Rectum* 1997; 40: 747-51.

Heald RJ, Moran BJ, Ryall RDH, Sexton R, MacFarlane JK. Rectal cancer. The Basingstoke experience of total mesorectal excision, 1978-1997. *Arch Surg* 1998; 133: 894-9.

Heldwein W, Dollhopf M, Rosch T, Meining A, Schmidtsdorff G, Hasford J et al. The Munich polypectomy study (MUPS): Prospective analysis of complications and risk factors in 4000 colonic snare polypectomies. *Endoscopy* 2005 Nov; 37 (11): 1116-22.

Hendren SK, Swallow CJ, Smith A, Lipa JE, Cohen Z, MacRae HM et al. Complications and sexual function after vaginectomy for anorectal tumors. *Dis Colon Rectum* 2007; 50: 810-6.

Henriksen MG, Hansen HV, Hesselø I (2002). Early oral nutrition after elective colorectal surgery: influence of balanced analgesia and enforced mobilization. *Nutrition* 18: 263-267.

Herbertson RA, Lee ST, Tebbutt N, Scott AM. The expanding role of PET technology in the management of patients with colorectal cancer. *Ann Oncol* 2007; 18: 1774-1781.

Heriot AG, Byrne CM, Lee P, Dobbs B, Tilney H, Solomon MJ et al. Extended Radical Resection: The Choice of Locally Recurrent Rectal Cancer. *Dis Colon Rectum* 2008; 51: 284-91.

Hermanek P, Gall FP. Early (microinvasive) colorectal carcinoma. Pathology, diagnosis, surgical treatment. *Int J Colorectal Dis* 1986; 1: 79-84.

Hershman MJ, Myint AS, Makin CA. Multi-modality approach in curative local treatment of early rectal carcinomas. *Colorectal Dis* 2003; 5: 445-50.

Hewitson P, Glasziou P, Irwig L, Towler B, Watson E. Screening for colorectal cancer using the faecal occult blood test, Hemoccult. *Cochrane Database of Systematic Reviews* 2006, Issue 4. Art. No.: CD001216.

Hewitt M, Petiti D, red. Interpreting the volume-outcome relationship in the context of cancer care. Washington, DC. The National Academies Press, 2001.

Hewitt M. Interpreting the volume-outcome relationship in the context of health care quality. Report from the Institute of Medicine, Washington, DC, 2000.

Hida J, Yasutomi M, Fujimoto K, Okuno K, Ieda S, Machidera N et al. Functional outcome after low anterior resection with low anastomosis for rectal cancer using the colonic J-pouch. Prospective randomised study for determination of optimum pouch size. *Dis Colon Rectum* 1996; 39: 986-91.

Hill GE, Grawley WH, Griffith KE, Forestner JE, Minei JP. Allogeneic blood transfusion increases the risk of postoperative bacterial infection: a meta-analysis. *J Trauma* 2003; 54: 908-14.

Ho YH, Tan M, Seow-Choen F. Prospective randomized controlled study of clinical function and anorectal physiology after low anterior resection: comparison of straight and colonic J pouch anastomoses. *Br J Surg* 1996; 83: 978-80.

Hoff PM, Ansari R, Batist G, Cox J, Kocha W, Kuperminc M et al. Comparison of oral capecitabine versus intravenous fluorouracil plus leucovorin as first-line treatment in 605 patients with metastatic colorectal cancer: results of a randomized phase III study. *J Clin Oncol* 2001 April 15; 19 (8): 2282-92.

Hohenberger W, Merkel S, Matzel K, Bittorf B, Papadopoulos T, Göhl J. The influence of abdomino-perianal (intersphincteric) resection of lower third rectal carcinoma on the rates of sphincter preservation and locoregional recurrence. *Colorectal Dis* 2006; 8: 23-33.

Hohenberger W, Matzel KE, Merkel S, Papadopoulos T, Weber K. Standardised Surgery for Colonic Cancer: Complete mesocolic excision (CME) and central ligation – Technical notes and outcome. *Colorectal Dis* 2008, Nov. 7.

Holm T, Ljung A, Häggmark T, Jurell G, Lagergren J. Extended abdominoperineal resection with gluteus maximus flap reconstruction of the pelvic floor for rectal cancer. *Br J Surg* 2007; 94: 232-8.

Holte K, Foss N, Andersen J, Valentiner L, Lund C, Bie P et al (2007). Liberal or restrictive fluid administration in fast-track colonic surgery: a randomized, double-blind study. *Br J Anaesth* 99: 500-508.

Holte K, Kehlet H (2006). Fluid therapy and surgical outcomes in elective surgery: a need for reassessment in fast-track surgery. *J Am Coll Surg* 2002: 971-989.

Huebner RH, Park KC, Shepherd JE et al. A meta-analysis of the literature for whole-body FDG PET detection of recurrent colorectal cancer. *J Nucl Med* 2000; 41: 1177-1189.

Hughes R, Glynne-Jones R, Grainge J et al. Can pathological complete response in the primary tumour following preoperative pelvic chemoradiotherapy for T3eT4 rectal cancer predict for sterilisation of pelvic lymph nodes, a low risk of local recurrence and the appropriateness of local excision? *Int J Colorectal Dis* 2006; 21 (1): 11-17.

Hunerbein M, Below C, Schlag PM. Three-dimensional endorectal ultrasonography for staging of obstructing rectal cancer. *Dis Colon Rectum* 1996; 39: 636-41.

Hunerbein M, Pegios W, Rau B, Vogl TJ, Felix R, Schlag PM. Prospective comparison of endorectal ultrasound, three-dimensional endorectal ultrasound, and endorectal MRI in the preoperative evaluation of rectal tumors. Preliminary results. *Surg Endosc* 2000; 14: 1005-1009.

Hunerbein M, Totkas S, Moesta KT, Ulmer C, Handke T, Schlag PM. The role of transrectal ultrasound-guided biopsy in the postoperative follow-up of patients with rectal cancer. *Surgery* 2001; 129: 164-169.

Imperiale TF, Ransohoff DF, Itzkowitz SH, Turnbull BA, Ross ME. Colorectal Cancer Study Group. Fecal DNA versus fecal occult blood for colorectal cancer screening in an average-risk population. *N Engl J Med* 2004; 351: 2704-14.

Improved survival with preoperative chemoradiotherapy for respectable rectal cancer. Swedish Rectal Cancer trial. *NEJM* 1997; 336: 980-7.

Irvin TT, Greaney MG. The treatment of colonic cancer presenting with intestinal obstruction. *Br J Surg* 1977; 64: 741-4.

Iversen L. Kvaliteten af den kirurgiske behandling for colorektal cancer relateret til antallet af udførte operationer og kirurgisk uddannelse. En litteraturgennemgang. Upubliceret.

Iversen LH, Harling H, Laurberg S, Wille-Jørgensen P, Danish Colorectal Cancer Group. Influence of caseload and surgical speciality on outcome following surgery for colorectal cancer: a review of evidence. Part 1: Short-term outcome. *Colorectal Dis* 2007; 9: 28-37.

Iversen LH, Harling H, Laurberg S, Wille-Jørgensen P, Danish Colorectal Cancer Group. Influence of caseload and surgical speciality on outcome following surgery for colorectal cancer: a review of evidence. Part 2: long-term outcome. *Colorectal Dis* 2007; 9: 38-46.

Jagoditsch M, Lisborg PH, Jatzko GR, Wette V, Kropfisch G, Denk H et al. Long-term prognosis for colon cancer related to consistent radical surgery: Multivariate analysis of clinical, surgical and pathologic variables. *World J Surg* 2000; 24: 1264-70.

Jain A, Falzarano J, Jain A, Decker R, Okubo G, Fujiwara D. Outcome of 5,000 flexible sigmoidoscopies done by nurse endoscopists for colorectal cancer screening in asymptomatic patients. *Hawaii Med J* 2002; 61: 118-20.

Jakobsen D, Sonne E, Andreasen J, Kehlet H (2006). Convalescence after colonic surgery with fast-track vs conventional care. *Colorectal Dis* 8: 683-687.

Jakobsen DH, Hallin M, Kehlet H (1999). Nye plejeregninger efter colon kirurgi. *Sygeplejersken* 46: 37-39.

Janjan NA, Abbruzzese J, Pazdur R et al. Prognostic implications of response to preoperative infusional chemoradiation in locally advanced rectal cancer. *Radiother Oncol* 1999; 51 (2): 153-160.

Jass JR. Towards a molecular classification of colorectal cancer. *Int J Colorecti Dis* 1999; 14: 194-200.

Jayne DG, Guillou PJ, Thorpe H, Quirke P, Copeland J, Smith AM et al. Randomized trial of laparoscopic-assisted resection of colorectal carcinoma: 3-years results of the UK MRC CLASICC Trial Group. *J Clin Oncol* 2007; 25: 3061-3068.

Jeffery GM, Hickey BE, Hider P. Follow-up strategies for patients treated for non-metastatic colorectal cancer. *Cochrane Database Syst Rev*. 2002; (1): CD002200.

Jeffery M, Hickey BE, Hider PN. Follow-up strategies for patients treated for non-metastatic colorectal cancer. *Cochrane Database of Systematic Reviews* 2007; issue 1.

Jenkins MA, Hayashi S, O'Shea AM, Burgart LJ, Smyrk TC, Shimizu D et al. Pathology features in Bethesda guidelines predict colorectal cancer microsatellite instability: a population-based study. *Gastroenterology* 2007; 133: 48-56.

Jensen LH, Lindebjerg J, Byriel L, Kolvraa S, Cruger DG. Strategy in clinical practice for classification of unselected colorectal tumours based on mismatch repair deficiency. *Colorectal Dis* 2008; 10: 490-497.

Jensen LS, Puho E, Pedersen L, Mortensen FV, Sørensen HT. Long-term survival rate of patients transfused with leucocyte-depleted and buffy-coat-poor blood during colorectal surgery – secondary publication. I An eight-year post-trial follow-up study. *Ugeskr Laeger* 2006 Jan 30; 168 (5): 481-4.

Jess P, Christiansen J, Bech P. Quality of life after anterior resection versus abdominoperineal extirpation for rectal cancer. *Scand J Gastroenterol* 2002; 37: 1201-4.

Jestin P, Heurgren M, Pålman L, Glimelius B, Gunnarsson U. Elective surgery for colorectal cancer in a defined Swedish population. *Eur J Surg Oncol* 2004; 30: 26-33.

Jestin P, Nilsson J, Heurgren M et al. Emergency surgery for colonic cancer in a defined population. *Br J Surg* 2005; 92 (1): 94-100.

Jestin P, Pålman L, Gunnarsson U. Risk factors for anastomotic leakage after rectal cancer surgery: a case-control study. *Colorectal Dis* 2008; 10: 715-21.

Jesus EC, Karliczek A, Matos D, Castro AA, Atallah An. Prophylactic anastomotic drainage for colorectal surgery. *Cochrane Database Syst Rev* 2004; 4: CD002100.

Johnson DS, Wong JH. The impact on nodal staging of lymphatic mapping in carcinoma of the colon and rectum. *Semin Oncol* 2004; 31: 403-8.

Johnson CD, Chen MH, Toledano AY, Keiken JP, Dachman A, Kuo MD et al. Accuracy of CT colonography for detection of large adenomas and cancers. *N Engl J Med* 2008; 359: 1207-17.

Jones OM, Rees M, John TG, Bygrave S, Plant G. Biopsy of resectable colorectal liver metastases causes tumour dissemination and adversely affects survival after liver resection. *Br J Surg* 2005; 92: 1165-1168.

Jonker DJ, O'Callaghan CJ, Karapetis CS, Zalberg JR, Tu D, Au HJ et al. Cetuximab for the treatment of colorectal cancer. *N Engl J Med* 2007 November 15; 357 (20): 2040-8.

Jung B, Pahlman L, Nyström PO, Nilsson E (2007). Multicentre randomized clinical trial of mechanical bowel preparation in elective colonic resection. *Br J Surg* 94: 689-695.

Järvinen HJ, Aarnio M, Mustonen H, Aktan-collan K, Aaltonen LA, Peltomaki P et al. Controlled 15-year trial on screening for colorectal cancer in families with hereditary nonpolyposis colorectal cancer. *Gastroenterology* 2000; 118: 829-834.

Jørgensen OD, Kronborg O, Fenger C. A randomised study of screening for colorectal cancer using faecal occult blood testing: Results after 13 years and seven biennial screening rounds. *Gut* 2002; 50: 29-32.

Jørgensen LN, Lausen I, Rasmussen MS, Wille-Jørgensen P, Bergqvist D. Prolonged thromboprophylaxis with low molecular weight heparin (tinzaparin) following major general surgery. *Blood* 2003; 102: 1168.

Kakkar W, Cohen AT, Edmonson RA, Phillips MJ, Cooper DJ, Das SK et al. Low molecular weight versus standard heparin for prevention of venous thromboembolism after major abdominal surgery. The Thromboprophylaxis Collaborative Group. *Lancet* 1993; 341: 259-265.

Kalady MF, Mantyh CR, Petrofski J, Ludwig KA. Routine contrast imaging of low pelvic anastomosis prior to closure of defunctioning ileostomy: is it necessary? *J Gastrointest Surg* 2008; 12: 1227-31.

Kalofonos HP, Aravantinos G, Kosmidis P, Papakostas P, Economopoulos T, Dimopoulos M et al. Irinotecan or oxaliplatin combined with leucovorin and 5-fluorouracil as first-line treatment in advanced colorectal cancer: a multicenter, randomized, phase II study. *Ann Oncol* 2005 June; 16 (6): 869-77.

Kanehira E, Yamashita Y, Omura K, Kinoshita T, Kawakami K, Watan G. Early clinical results of endorectal surgery using a newly designed rectal tube with a side window. *Surg Endosc* 2002; 16: 14-17.

Kapiteijn E, Marijnen CA, Nagtegaal ID, Putter H, Steup WH, Wiggers T et al. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer. *N Engl J Med* 2001; 345: 638-646.

Kapiteijn E, Putter H, van de Velde CJ, Cooperative investigators of the Dutch ColoRectal Cancer Group. Impact of the introduction and training of total mesorectal excision on recurrence and survival in rectal cancer in The Netherlands. *Br J Surg* 2002; 89: 1142-9. Karanjia 1994.

Karp I, Behloul H, Leloir J, Pilote L. Statins and cancer risk. *Am J Med* 2008; 121: 302-9.

Katballe N. Hereditary Non-Polyposis Colorectal Cancer in West Denmark. Prospective Polpulation-based studies. PhD thesis 2001; 1-104.

Kehlet H (2008). Fast-track colorectal surgery. *Lancet* 371: 791-793.

Kewenter J, Björk S, Haglind E, Smith L, Svanvik J, Ahrén C. Screening and rescreening for colorectal cancer. A controlled trial of fecal occult blood testing in 27,700 subjects. *Cancer* 1988; 62: 645-51.

Keymling M. Colorectal Stenting. *Endoscopy* 2003; 35: 234-238.

Khan AA, Wheeler JM, Cunningham C, George B, Kettlewell M, Mortensen NJ. The management and outcome of anastomotic leaks in colorectal surgery. *Colorectal Dis* 2007 Dec 7 (Epub ahead of print).

Khatri VP, Rodriguez-Bigas MA, Petrelli NJ. Perineal dissection of synchronous abdominoperineal resection of the rectum: An anatomical description. *Arch Surg* 2003; 138: 553-9.

Khoo CK, Vickery CJ, Forsyth N, Vinall NS, Eyre-Brook IA. A prospective randomized controlled trial of multimodal perioperative management protocol in patients undergoing elective colorectal resection for cancer. *Ann Surg* 2007; 245: 867-72.

Khot UP, lang AW, Murali K et al. Systemic review of the efficacy and safety of colorectal stents. *Br J Surg* 2002; 89 (9): 1016-

Kiesslich R, Jung M, DiSario JA, Galle PR, Neurath MF. Perspectives of chromo and magnifying endoscopy. How, how much, when, and whom should we stain? *J Clin Gastroenterol* 2004; 38: 7-13.

Kievit J. Follow-up of patients with colorectal cancer: Numbers needed to test and treat. *Eur J Cancer* 2002; 38: 986-99.

Kievit J. Colorectal cancer follow-up: a reassessment of empirical evidence on effectiveness. *Eur J Surg Oncol* 2000; 26: 322-328.

Kikuchi R, Takano M, Takagi K, Fujimoto N, Nozaki R, Fujuyoshi T et al. Management of early invasive colorectal cancer. Risk of recurrence and clinical guidelines. *Dis Colon Rectum* 1995; 38: 1286-95.

Kim CJ, Yeatman T, Coppola D, Trotti A, Williams B, Barthel JS et al. Local excision of T2 and T3 rectal cancers after downstaging chemoradiation. *Ann Surg* 2001; 3: 352-59.

Kim HS, Kim TI, Kim WH, Kim YH, Yang SK et al. Risk factors for immediate postpolypectomy bleeding of the colon: A multicenter study. *Am J Gastroenterol* 2006 Jun; 101 (6): 1333-41.

Kim SH, Milsom JW, Church JM, Ludwig KA, Garcia-Ruiz A, Okuda J et al. Perioperative tumor localization for laparoscopic colorectal surgery. *Surg Endosc* 1997; 11: 1013-1016.

Kim YK, Ko SW, Hwang SB, Kim CS, Yu HC. Detection and characterization of liver metastases: 16-slice multidetector computed tomography versus superparamagnetic iron oxide-enhanced magnetic resonance imaging. *Eur Radiol* 2006; 16: 1337-1345.

Kimmey MB. Endoscopic methods (other than stents) for palliation of rectal carcinoma. *J Gastrointest Surg* 2004; 8: 270-3.

King PM, Blazeby JM, Ewings P, Franks PJ, Longman RJ, Kendrick AH et al. Randomized clinical trial comparing laparoscopic and open surgery for colorectal cancer within an enhanced recovery programme. *Br J Surg* 2006; 93: 300-308.

Kinkel K, Lu Y, Both M, Warren RS, Thoeni RF. Detection of hepatic metastases from cancers of the gastrointestinal tract by using non-invasive imaging methods (US, CT, MR imaging, PET): a meta-analysis. *Radiology* 2002; 224: 748-756.

Kirkland KB, Briggs JP, Trivette SL, Wilkinson WE, Sexton DJ. The impact of surgical-site infections in the 1990s: attributable mortality, excess length of hospitalization, and extra costs. *Infect Control Hosp Epidemiol* 1999; 20: 725-30.

Kjeldsen BJ, Kronborg O, Fenger C, Jørgensen OD. A prospective randomized study of follow-up after radical surgery for colorectal cancer. *Br J Surg* 1997a; 84: 666-669.

Kjeldsen BJ, Kronborg O, Fenger C, Jørgensen OD. The patterns of recurrent colorectal cancer in a prospective randomised study and the characteristics of diagnostic tests. *Int J Colorect Dis* 1997b; 12: 329-334.

Kjeldsen BJ, Thorsen H, Whalley D, Kronborg O. Influence of follow-up on health-related quality of life after radical surgery for colorectal cancer. *Scand J Gastroenterol* 1999; 34: 509-515.

Kjeldsen BJ. Evaluation of long-term follow-up in patients having had surgery for colorectal cancer with curative intent. Ph.D. thesis 1998. Odense, Denmark, Faculty of Health Science, Odense University, Denmark.

Knowles G, Sherwood L, Dunlop MG, Dean G, Jodrell, McLean C et al. Develoeping and piloting a nurse-led model of follow-up in the multidisciplinary management of colorectal cancer. *Eur J Oncol Nurs* 2007; 11: 121-223.

Ko CW, Dominitz JA, Nguyen TD. Fecal occult blood testing in a general medical clinic: comparison between guaiac-based and immunochemical-based test. *Am J Med* 2003; 115: 111-4.

Koh DM, Brown G, Temple L, Raja A, Toomey P, Bett N et al. Rectal cancer: mesorectal lymph nodes at MR imaging with USPIO versus histopathologic findings – initial observations. *Radiol* 2004; 231: 91-9.

Koh DM, Chau I, Tait D, Wotherspoon A, Cunningham D, Brown G. Evaluating mesorectal lymph nodes in rectal cancer before and after neoadjuvant chemoradiation using thin-section T2-weighted magnetic resonance imaging. *Int J Radiat Oncol Biol Phys*. 2008; 71: 456-61.

Kohne CH, Wils J, Lorenz M, Schoffski P, Voigtmann R, Bokemeyer C et al. Randomized phase III study of high-dose fluorouracil given as a weekly 24-hour infusion with or without leucovorin in advanced colorectal cancer: European organization of Research and Treatment of Cancer Gastrointestinal Group Study 40952. *J Clin Oncol* 2003 October 15; 21 (20): 3721-8.

Kondrup J, Rasmussen HH, Hamberg O, Stanga Z; Ad Hoc ESPEN Working Group. Nutritional risk screening (NRS 2002): a new method based on an analysis of controlled clinical trials. *Clin Nutr* 2003; 22: 321-36.

Konishi F, Muto T, Kanazawa K, Morioka Y. Intraoperative irrigation and primary resection for obstructing lesions of the left colon. *Int. J Colorectal Dis* 1988; 3: 204-6.

Konopke R, Bunk A, Kersting S. The role of contrast-enhanced ultrasound for focal liver lesion detection: an overview. *Ultrasound Med Biol* 2007; 33: 1515-1526.

Konopke R, Kersting S, Bergert H et al. Contrast-enhanced ultrasonography to detect liver metastases: a prospective trial to compare transcutaneous unenhanced and contrast-enhanced ultrasonography in patients undergoing laparotomy. *Int J Colorectal Dis* 2007; 22: 201-207.

Koopman M, Antonini NF, Douma J, Wals J, Honkoop AH, Erdkamp FL et al. Sequential versus combination chemotherapy with capecitabine, irinotecan, and oxaliplatin in advanced colorectal cancer (CAIRO): a phase III randomised controlled trial. *Lancet* 2007 July 14; 370 (9582): 135-42.

Koshiji M, Yonekura Y, Saito T, Sakaida N, Uemura Y, Yoshioka K. Genetic alterations in normal epithelium of colorectal cancer patients may be a useful indicator for subsequent metachronous tumor development. *Ann Surg Oncol* 2002; 9: 580-6.

Kristiansen C, Loft A, Berthelsen AK, Graff J, Lindebjerg J, Bisgaard C et al. PET/CT and histopathologic response to preoperative chemoradiation therapy in locally advanced rectal cancer. *Dis Colon Rectum* 2008; 51: 21-5.

Kronawitter U, Kemeny NE, Heelan R, Fata F, Fong Y. Evaluation of chech computed tomography in the staging of patients with potentially respectable liver metastases from colorectal carcinoma. *Cancer* 1999; 86: 229-235.

Kronborg O, Fenger C, Deichgräber E. Kolonoskopi efter radikal kirurgi for kolorektalcancer. Ti års prospektiv undersøgelse af 309 patienter. *Ugeskr Laeger* 1991; 153: 503-506.

Kronborg O, Fenger C, Olsen J, Bech K, Sondergaard O. Repeated screening for colorectal cancer with fecal occult blood test. A prospective randomized study at Funen, Denmark. *Scand J Gastroenterol* 1989; 24: 599-606.

Kronborg O, Fenger C, Olsen J, Jorgensen OD, Sondergaard O. Randomised study of screening for colorectla cancer with faecal-occult-blood test. *Lancet* 1996; 348: 1467-71.

Kronborg O, Fenger C, Sondergaard O, Pedersen KM, Olsen J. Initial mass screening for colorectal cancer with fecal occult blood test. A prospective randomized study at Funen in Denmark. *Scand J Gastroenterol* 1987; 22: 677-86.

Kronborg O, Regula J. Population screening for colorectal cancer: advantages and drawbacks. *Dig Dis* 2007; 25 (3): 270-3.

Kræftplan II. Sundhedsstyrelsens anbefalinger til forbedringer af indsatsen på kræftområdet. Sundhedsstyrelsen, 2005.

Kuebler JP, Wieand HS, O'Connell MJ, Smith RE, Colangelo LH, Yothers G et al. "Oxaliplatin combined with weekly bolus fluorouracil and leucovorin as surgical adjuvant chemotherapy for stage II and III colon cancer: results from NSABP C-07". *J Clin Oncol*, Vol. 25, No. 16, 2007, pp. 2198-2204.

Kuhry E, Bonjer HJ, Haglind E, Hop WC, Veldkamp R, Cuesta MA et al. Impact of hospital case volume on short-term outcome after laparoscopic operation for colonic cancer. *Surg Endosc* 2005; 19: 687-692.

Kuo LJ, LeuSY, Liu MC, Jian JJ, Hongiun CS, Chen CM. How aggressive should we be in patients with stage IV colorectal cancer? *Dis Colon Rectum* 2003; 46: 1646-52.

Kuvshinoff B, Maghfoor I, Miedema B, Bryer M, Westgate S, Wilkes J et al. Distal margin requirements after preoperative chemoradiotherapy for distal rectal carcinomas: are < or = 1 cm distal margins sufficient? *Ann Surg Oncol* 2001; 8: 163-9.

Labianca R, Floriani I, Cortesi E, Isa L, Zaniboni A, Marangolo M et al. Alternating versus continuous "FOLFIRI" in advanced colorectal cancer (ACC): A randomized "GISCAD" trial. *Proc Am Soc Clin Oncol* 2006; #3505.

Lacy AM, Garcia-Valdecasas JC, Delgado S, Castells A, Taura P, Pique JM et al. Laparoscopy-assisted colectomy versus open colectomy for treatment of non-metastatic colon cancer: a randomised trial. *Lancet* 2002; 359: 2224-2229.

Lahaye MJ et al. Imaging for predicting the risk factors – the circumferential resection margin and nodal disease – of local recurrence in rectal cancer: a meta-analysis. *Semin Ultrasound CT MR* 2005 Aug; 26 (4): 259-68.

Landmann RG, Wong WD, Hoepfl J, Shia J, Guillem JF, Temple LK et al. Limitations of early rectal cancer nodal staging may explain failure after local excision. *Dis Colon Rectum* 2007; 50: 1520-5.

Lange MM, den Dulk M, Bossema ER, Maas CP, Peeters KC, Rutten HJ et al. Risk factors for faecal incontinence after rectal cancer treatment. *Br J Surg* 2007; 94: 1278-84.

Lange MM, Buunen M, van de Velde CJ, Lange JF. Level of arterial ligation in rectal cancer surgery: low tie preferred over high tie. A Review. *Dis Colon Rectum* 2008; 51: 1139-45.

Larach SW, Patankar SK, Ferrara A, Williamson PR, Perozo SE, Lord AS. Complications of laparoscopic colorectal surgery. Analysis and comparison of early vs latter experience. *Dis Colon Rectum* 1997; 40: 592-596.

Larsen S, Wiig J, Emblemvaag H, Grøholt K, Hole K, Bentsen A et al. Extended TME in Locally Advanced Rectal Cancer (T4a) and the clinical role of MRI evaluated neo-adjuvant downstaging. *Colorectal Dis* 2008; Jul 25.

Larsen LPS, Rosenkilde M, Christensen H, Bang N, Bolvi L, Christensen T et al. The value of contrast enhanced ultrasonography in detection of liver metastases from colorectal cancer: A prospective double-blinded study. *Eur J Radiology* 2007; 62: 302-307.

Law WI, Chu KW, Ho JW, Chan CW. Risk factors for anastomotic leakage after low anterior resection with total mesorectal excision. *Am J Surg* 2000; 179: 92-6.

Law WL, Choi HK, Lee YM, Ho JW, Seto CL. Anastomotic leakage is associated with poor long-term outcome in patients after curative colorectal resection for malignancy. *Gastrointestinal Surg* 2007 Jan; 11 (1): 8-15.

Law WL, Chu KW. Abdominoperineal resection is associated with poor oncological outcome. *Br J Surg* 2004; 91: 1493-9.

Law WL, Chu KW. Anterior resection for rectal cancer with mesorectal excision: a prospective evaluation of 622 patients. *Ann Surg* 2004; 240: 260-8.

Law WL, Chus KW, Choi HK. Randomized clinical trial comparing loop ileostomy and loop transverse colostomy for faecal diversion following total mesorectal excision. *Br J Surg* 2002; 89: 704-8.

Law WL, Fan JK, Poon JT, Choi HK, Lo OS. Laparoscopic bowel resection in the setting of metastatic colorectal cancer. *Ann Surg Oncol* 2008; 15: 1424-1428.

Leaper M, Johnston MJ, Barclay M, Dobbs BR, Frizelle FA. Reasons for failure to diagnose colorectal carcinoma at colonoscopy. *Endoscopy* 2004; 36: 499-503.

Lee W, Lee D, Choi S, Chun H. Transanal endoscopic microsurgery and radical surgery for T1, T2 rectal cancers. *Surg Endosc* 2003; 17: 1283-87.

Lee MS. Update on techniques of colonic endoscopic mucosal resection. *Digestive Endoscopy* 2006; 18: 105-107.

Lejeune C, Bismuth MJ, Conroy T, Zanni C, Bey P, Bedenne L et al. Use for decision analyses model to assess the cost-effectiveness of 18F-FDG PET in the management of metachronous live metastases of colorectal cancer. *J Nucl Med* 2005; 46: 2020-2028.

Lembersky BC, Wieand HS, Petrelli NJ, O'Connell MJ, Colangelo LH, Smith RE et al. "Oral uracil and tegafur plus leucovorin compared with intravenous fluorouracil and leucovorin in stage II and III carcinoma of the colon: results from National Surgical Adjuvant Breast and Bowel Project Protocol C-061". *J Clin Oncol*, Vol. 24, No. 13, 2006, pp. 2059-2064.

Leung KL, Kwok SP, Lam SC, Lee JF, Yiu RY, Ng SS et al. Laparoscopic resection for rectosigmoid carcinoma: prospective randomised trial. *Lancet* 2004; 363: 1187-1192.

Levi Z, Rozen P, Hazazi R, Vilkin A, Waked A, Maoz E et al. A quantitative immunochemical fecal occult blood test for colorectal neoplasia. *Ann Intern Med* 2007; 146: 244-55.

Levin TR, Conell C, Shapiro JA, Chazan SG, Nadel MR, Selby JV. Complications of screening flexible sigmoidoscopy. *Gastroenterology* 2002; 123: 1786-92.

Levin B, Lieberman DA, McFarland B, Smith RA, Brooks D, Andrews KS et al. Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: a joint guideline from the American Cancer Society, the US Multi-Society Task Force on Colorectal Cancer, and the American College of Radiology. *CA Cancer J Clin* 2008; 58: 130-60.

Lewis SJ, Egger M, Sylvester PA, Thomas S (2001). Early enteral feeding versus "nil by mouth" after gastrointestinal surgery: systemic review and meta-analysis of controlled trials. *BMJ* 323: 773-776.

Lichtenstein P, Holm NV, Verkasalo PK, Iliadou A, Kaprio J, Koskenvuo M et al. Environmental and heritable factors in the causation of cancer – analyses of cohorts of twins from Sweden, Denmark, and Finland. *N Engl J Med* 2000; 343: 78-85.

Lievre A, Bachet JB et al. KRAS mutation as an independent prognostic factor in patients with advanced colorectal cancer treated with Cetuximab. *J Clin Oncol* 2008; 26: 374-9.

Linardou H, Dahabreh IJ, Kanaloupiti D, Siannis F, Bafaloukos D, Kosmidis P et al. Assessment of somatic k-RAS mutations as a mechanism associated with resistance to EGFR-targeted agents: a systematic review and meta-analysis of studies in advanced non-small-cell lung cancer and metastatic colorectal cancer. *Lancet Oncol* 2008; Preview before print.

Lindebjerg J, Spindler KL, Ploen J et al. The prognostic value of lymph node metastases and tumour regression grade in rectal cancer patients treated with long-course preoperative chemoradiotherapy. *Colorectal Dis* (in press).

Lindor NM, Rabe K, Petersen GM, Haile R, Casey G, Baron J et al. Lower cancer incidence in Amsterdam-I criteria families without mismatch repair deficiency: familial colorectal cancer type X. *Jama* 2005; 293: 1979-1985.

Lloyd GM, Sutton CD, Marshall LJ, Baragwanath P, Jameson JS, Scott AD. Transanal endoscopic microsurgery – lessons from a single UK centre series. *Colorectal Dis* 2002; 4: 467-72.

Lopez-Kostner F, Lavery IC, Hool GR, Rybicki LA, Fazio VW. Total mesorectal excision is not necessary for cancers of the upper rectum. *Surgery* 1998; 124: 612-7.

Lourenco T, Murray A, Grant A, McKinley A, Krukowski Z, Vale L. Laparoscopic surgery for colorectal cancer: safe and effective? – A systematic review. *Surg Endosc* 2008; 22: 1146-1160.

Luna-Pérez P, Rodriguez-Ramirez SE, De La Barrera MG, Zeferino M, Labastida S. Multivisceral Resection for Colon Cancer. *J Surg Oncol* 2002; 80: 100-04.

Lynch HAT, De La CA. Hereditary colorectal cancer. *N Engl J Med* 2003; 348: 919-932.

MacDonald JS. Clinical overview: adjuvant therapy of gastrointestinal cancer. *Cancer Chemother Pharmacol* 2004; 54 (Suppl 1): S4-11.

Macfie J, Woodcock NP, Palmer MD, Walker A, Townsend S, Mitchell CJ (2000). Oral dietary supplements in pre- and postoperative surgical patients: a prospective and randomized clinical trial. *Nutrition* 16: 723-728.

Machado M, Nygren J, Goldman S, Ljungqvist O. Functional and physiologic assessment of the colonic reservoir or side-to-end anastomosis after low anterior resection for rectal cancer: a two-year follow-up. *Dis Colon Rectum* 2005; 48: 29-36.

MacKay G, Fearon K, McConnachie A, Serpell M, Molloy R, O'Dwyer P (2006). Randomized clinical trial of the effect of postoperative intravenous fluid restriction on recovery after elective colorectal surgery. *Br J Surg* 93: 1469-1474.

Madsen MR, Harling H. Kontrol af Patienter efter radikal operation for kolorektalcancer. *Ugeskr Laeger* 2004.

Maessen J, Dejong C, Hausel J, Nygren J, Lassen K, Andersen J et al (2007). A protocol is not enough to implement an enhanced recovery programme for colorectal resection. *Br J Surg* 94: 224-231.

Maetani S, Onodera H, Nishikawa T. Long-Term Cure in Surgery for Extrarectal Pelvic Recurrence of Rectal cancer. *Dis Colon Rectum* 2007; 50: 1558-67.

Mahmood T, Darzi A. A study to validate the colonoscopy simulator. *Surg Endosc* 2003; 17: 1583-9.

Mainar A, De Gregorio Ariza MA, Tejero E et al. Acute colorectal obstruction: Treatment with self-expandable metallic stents before scheduled surgery – results of a multicenter study. *Radiology* 1999; 210: 65-69.

Maindrault-Goebel F, Lledo G, Chibaudel B, Mineur L, Andre T, Bennamoun M et al. Final results of OPTIMOX2, a large randomized phase II study of maintenance therapy or chemotherapy-free intervals (CFI) after FOLFOX in patients with metastatic colorectal cancer (MRC): A GERCOR study. *Proc Am Soc Clin Oncol* 2007; #4013.

Malmstrøm ML, Meisner S. Endoscopic mucosal resection of the colon and rectum. *Ugeskr Laeger* 2008; 170: 1738-9.

Mandard AM, Dalibard F et al. Pathologic assessment of tumour regression after preoperative chemoradiotherapy of esophageal carcinoma. *Cancer* 1994; 73: 2680-6.

Mandel JS, Bond JH, Church TR, Snover DC, Bradley GM, Schuman LM et al. Reducing mortality from colorectal cancer by screening for fecal occult blood. Minnesota Colon Cancer Control Study. *N Engl J Med* 1993; 328: 1365-71.

Mandel JS, Church TR, Bond JH, Ederer F, Geisser MS, Mongin SJ et al. The effect of fecal occult-blood screening on the incidence of colorectal cancer. *N Engl J Med* 2000; 343: 1603-7.

Mandel JS, Church TR, Ederer F, Bond JH. Colorectal cancer mortality: effectiveness of biennial screening for fecal occult blood. *J Natl Cancer Inst* 1999; 91: 434-7.

Marijnem CAM et al. Acute side effects and complications after short-term preoperative radiotherapy combined with total mesorectal excision in primary rectal cancer: report of a multicenter randomized trial. *J Clin Oncol* 2002 Feb 1; 20 (3): 817-25.

Marijnen CA, van de Velde CJ, Putter H, van den Brink M, Maas CP, Martijn H et al. Impact of short-term preoperative radiotherapy on health-related quality of life and sexual functioning in primary rectal cancer: report of a multicenter randomized trial. *J Clin Oncol* 2005; 23: 1847-58.

Markantonis SL, Kostopanoagiotou G, Panidis D, Smirniotis V, Voros D. 2004. Effects of blood loss and fluid volume replacement on serum and tissue gentamicin concentrations during colorectal surgery. *Clin Ther* 26: 271-281.

Marks JH, Kawun UB, Hamdan W, Marks G. Redefining contraindications to laparoscopic colorectal resection for high-risk patients. *Surg Endosc* 2008; 18 marts, Epub ahead of pring.

Marr R, Birbeck K, Garvican J, Macklin CP, Tiffin NJ, Parsons WJ et al. The modern abdominoperineal excision: the next challenge after total mesorectal excision. *Ann Surg* 2005; 242: 74-82.

Marsh PJ, James RD, Schofield PF. Adjuvant preoperative radiotherapy for locally advanced rectal carcinoma. Results of a prospective, randomized trial. *Dis Colon Rectum* 1994; 37: 1205-14.

Martinez-Santos C, Lobato RF, Fradejas JM et al. Self-expandable stent before elective surgery vs. emergency surgery for the treatment of malignant colorectal obstructions: Comparison of primary anastomosis and morbidity rates. *Dis Colon Rectum* 2002; 45: 401-406.

Martling AL, Holm T, Rutqvist L-E et al. Effect of a surgical training programme on outcome of rectal cancer in the county of Stockholm. *Lancet* 2000; 356: 93-6.

Martling AL, Holm T, Rutqvist LE et al. Effect of a surgical training programme on outcome of rectal cancer in the County of Stockholm. Stockholm Colorectal Cancer Study Group, Basingstoke Bowel Cancer Research Project. *Lancet* 2000; 356: 93-96.

Marusch F, Koch A, Schmidt U, Geibetaler S, Dralle H, Saeger HD et al. Value of a protective stoma in low anterior resections for rectal cancer. *Dis Colon Rectum* 2002; 45: 1164-71.

Maslekar S, Pillinger Sh, Monson JR. Transanal endoscopic microsurgery for carcinoma of the rectum. *Surg Endosc* 2007; 21: 97-102.

Matthiessen P, Halböök O, Rutegard J et al. Population-based study of risk factors for postoperative death after anterior resection of the rectum. *Br J Surg* 2006; 93: 498-503.

Matthiessen P, Hallböök O, Andersson M et al. Risk factors for anastomotic leakage after anterior resection of the rectum. *Colorectal Dis* 2004; 6: 462-9.

Matthiessen P, Hallböök O, Rutegård J, Simert G, Sjö Dahl R. Defunctioning stoma reduces symptomatic anastomotic leakage after low anterior resection of the rectum for cancer: a randomized multicenter trial. *Ann Surg* 2007; 246: 207-14.

Matthiessen P, Henriksson M, Hallböök O, Grunditz E, Norén B, Arbmán G. Increase of serum C-reactive protein is an early indicator of subsequent symptomatic anastomotic leakage after anterior resection. *Colorectal Dis* 2008; 10: 75-80.

Maurer CA. Urinary and sexual function after total mesorectal excision. *Recent Results Cancer Res* 2005; 165: 196-204.

McAnena OJ, Heald RJ, Lockhart-Mummery HE. Operative and functional results of total mesorectal excision with ultra-low anterior resection in the management of carcinoma of the lower one-third of the rectum. *Surg Gynecol Obstet* 1990; 170: 517-21.

McArdle CS and Hole DS. Emergency presentation of colorectal cancer is associated with poor 5-year survival. *Br J Surg* 2004; 91 (5): 610-17.

McArdle CS, hole D, Hansell D, Blumgart LH, Wood CB. Prospective study of colorectal cancer in the west of Scotland: 10 year follow-up. *Br J Surg* 1990; 77: 280-2.

McArdle CS, Hole D. Impact of variability among surgeons on postoperative morbidity and mortality and ultimate survival. *BMJ* 1991; 302: 1501-5.

McArdle CS, McMillan DC, Hole DJ. Impact of anastomotic leakage on long-term survival of patients undergoing curative resection for colorectal cancer. *Br J Surg* 2005; 92: 1150-4. McFarlane 1993.

Meisner S, Hensler M, Knop FK, West F, Wille-Jørgensen P. Self-expanding metal stents for colonic obstruction: Experience from 104 procedures in a single center. *Dis Colon Rectum* 2004; 47: 444-50.

Mella J, Datta SN, Biffin A, Radcliffe AG, Steele RJ, Stamatakis JD. Surgeons' follow-up practice after resection of colorectal cancer. *Ann R Coll Surg Engl* 1997; 79: 206-209.

Mentges B, Buess G, Effinger G, Mannicke K, Becker HD. Indications and results of local treatment of rectal cancer. *Br J Surg* 1997; 84: 348-51.

Mercury Study Group. Diagnostic accuracy of preoperative magnetic resonance imaging in predicting curative resection of rectal cancer: prospective observational study. *BMJ* 2006; 14: 333-779.

Merkel S, Mansmann U, Papadopoulos T, Wittekind C, Hohenberger W, Hermanek P. "The prognostic inhomogeneity of colorectal carcinomas Stage III: a proposal for subdivision of Stage III". *Cancer*, Vol. 92, No. 11, 2001, pp. 2754-2759.

Meyerhardt JA, Mayer RJ. Systemic therapy for colorectal cancer. *N Engl J Med* 2005 February 3; 352 (5): 476-87.

Michelassi F, Vannucci L, Ayala JJ, Chappel R, Goldberg R, Block GE. Local recurrence after curative resection of colorectal adenocarcinoma. *Surgery* 1990; 108: 787-93.

Miettinen RP, Laitinen ST, Mäkela JT, Pääkkönen ME (2000). Bowel preparation with oral polyethylene glycol electrolyte solution vs. no preparation in elective open colorectal surgery: prospective, randomized study. *Dis Colon Rectum* 43: 669-675.

Mismetti P, Laporte S, Darmon JY, Buchmuller A, Decousus H. Meta-analysis of low molecular weight heparin in the prevention of venous thromboembolism in general surgery. *Br J Surg* 2001; 88: 913-930.

Mitry E et al. Adjuvant chemotherapy after potentially curative resection of metastases from colorectal cancer: a pooled analysis of two randomized trials. *J Clin Oncol* 2008; 26: 4906-11.

Moertel CG, Fleming TR, Macdonald JS, Haller DG, Laurie JA, Goodman P et al. "Levamisole and fluorouracil for adjuvant therapy of resected colon carcinoma". *N Engl J Med*, Vol. 322, No. 6, 1990, pp. 352-358.

Mohiuddin M, Marks J, Marks G. Management of rectal cancer: short – vs long-course preoperative radiation. *Int J Radiat Oncol Biol Phys* 2008; 72: 636-43.

Mohiuddin N, Hayne M, Regine W et al. Prognostic significance of post-chemoradiation stage following preoperative chemotherapy and radiation for advanced/recurrent rectal cancers. *Int J Radiat Oncol Biol Phys* 2000; 48 (4): 1075-1080.

Moore HG, Riedel E, Minsky BD, Saltz L et al. Adequacy of 1-cm distal margin after restorative rectal cancer resection with sharp mesorectal excision and preoperative combined-modality therapy. *Ann Surg Oncol* 2003; 10: 80-5.

Morikawa E, Yasutomi M, Shindou K, Matsuda T, Mori N, Hida J et al. Distribution of metastatic lymph nodes in colorectal cancer by the modified clearing method. *Dis Colon Rectum* 1994; 37: 219-23.

Morris EJA, Maughan NJ, Forman D, Quircke P. Who to treat with adjuvant therapy in Dukes B/stage II colorectal cancer? The need for high quality pathology. *Gut* 2007; 56: 1419-1425.

Mortensen N. Rectal and anal endosonography. *Gut* 1992; 33: 148.

Mulhall BP, Veerappan GR, Jackson JL. Meta-analysis: computed tomographic colonography. *Ann Intern Med* 2005; 142: 635-50.

Murray JJ, Schoetz DJ Jr., Collier JA, Roberts PL, Veidenheimer MC. Intraoperative colonic lavage and primary anastomosis in nonelective colon resection. *Dis Colon Rectum* 1991; 34: 527-31.

Mynster T, Christensen IB, Moesgaard F, Nielsen HJ, The RANX05 Colorectal Cancer Study Group. The effects of the combination of blood transfusion and postoperative infectious complications on prognosis after colorectal cancer surgery. *Br J Surg* 2000; 87: 1553-62.

Mynster T, Nielsen HJ, Harling H, Bülow S; Danish TME-group, RANX05-group. Blood loss and transfusion after total mesorectal excision and conventional rectal cancer surgery. *Colorectal Dis* 2004 Nov; 6 (6): 452-7.

Myrhøj T, Andersen MB, Bernstein I. Screening for urinary tract cancer with urine cytology in Lynch syndrome and familial colorectal cancer. *Fam Cancer* 2008.

Myrhøj T, Bisgaard ML, Bernstein I, Svendsen LB, Søndergaard JO, Bülow S. Hereditary non-polyposis colorectal cancer: Clinical features and survival: results from the Danish HNPCC-register. *Scand J Gastroenterol* 1997; 32: 572-576.

Mäkelä JT, Kiviniemi H, Laitinen S. Risk factors for anastomotic leakage after left-sided colorectal resection with rectal anastomosis. *Dis Colon Rectum* 2003; 46: 653-60.

Mäkelä JT, Seppo OL, Kairaluoma MI. Five-year follow-up after radical surgery for colorectal cancer. *Archives of Surgery* 1995; 130: 1062-1067.

Nagtegaal ID, Marijnen CA, Kranenbarg EK, van de Velde CM, van Krieken JH. Pathology Review Committee; Cooperative Clinical Investigators. Circumferential margin involvement is still an important predictor of local recurrence in rectal carcinoma: not one millimetre but two millimetres is the limit. *Am J Surg Pathol* 2002; 26: 350-7.

Nagtegaal ID, Quirke P. What is the role for the circumferential margin in the modern treatment of rectal cancer? *J Clin Oncol* 2008; 26: 303-12.

Nagtegaal ID, van de Velde CJ, Marijnen CA, van Krieken JH, Quirke P, Dutch Colorectal Cancer Group et al. Low rectal cancer: a call for a change of approach in abdominoperineal resection. *J Clin Oncol* 2005; 23: 9257-64.

Nascimbeni R, Burgart LJ, Nivatvongs S, Larson DR. Risk of lymph node metastasis in T1 carcinoma of the colon and rectum. *Dis Colon Rectum* 2002; 45: 200-6.

Nasu K, Kuroki Y, Nawano S et al. Hepatic metastases: diffusion-weighted sensitivity encoding versus SPIO-enhanced MR imaging. *Radiology* 2006; 239: 122-130.

Nelson H, Petrelli N, Carlin A, Couture J, Fleshman J, Guillem J et al. Guidelines 2000 for Colon and Rectal Cancer Surgery. *J Natl Cancer Inst* 2001; 93: 583-96.

Nelson R, Edwards S, Tse B (2005). Prophylactic nasogastric decompression after abdominal surgery. *Cochrane Database Syst Rev* CD004929.

Nesbakken A, Nygaard K, Westerheim O, Mala T, Lunde OC. Local recurrence after mesorectal excision for rectal cancer. *Eur J Surg Oncol* 2002; 28: 126-34.

Nesbakken A, Nygaard K, Lunde OC et al. Anastomotic leak following mesorectal excision for rectal cancer: true incidence and diagnostic challenges. *Colorectal Dis* 2005; 7: 576-81.

Nespli A, Gianotti L, Totis M, Bovo G, Nespoli L, Chiodini P et al. 2004. Correlation between postoperative infections and long-term survival after colorectal resection for cancer. *Tumori* 90: 485-490.

Neugut AI, Lautenbach E, Abi-Rached B, Forde KA. Incidence of adenomas after curative resection for colorectal cancer. *AM J Gastroenterol* 1996; 91: 2096-2098.

NHS National Institute for Health and Clinical Excellence. Laparoscopic surgery for colorectal cancer. Review of NICE technology appraisal 17. National Institute for Health and Clinical Excellence. 2006.

Nicholson FB, Barro JL, Atkin W, Lilford R, Patrick J, Williams CB et al. Review article: population screening for colorectal cancer. *Aliment Pharmacol Ther* 2005; 22: 1069-1077.

Nickelsen TN, Jørgensen T, Kronborg O. Lifestyle and 30-day complications to surgery for colorectal cancer. *Acta Oncol* 2005; 44: 218-23.

Nielsen MB. Anorektal UL-scanning. *Ugeskr Laeger* 2001; 163: 4377-9.

Nielsen JD, Rasmussen HM, Husted S. The antithrombotictreated patient. *Ugeskr Lager* 2006; 168: 4296-4299.

Niël-Weise BS, van den Broek PJ (2005). Urinary catheter policies for short-term bladder drainage in adults. *Cochrane Database Syst Rev Issue 3:CD004203*.

Nilbert M, Wikman FP, Hansen TV, Krarup HB, Orntoft TF, Nielsen FC et al. Major contribution from recurrent alterations and MSH6 mutations in the Danish Lynch syndrome population. *Fam Cancer* 2008.

Nivatvongs S. Surgical management of early colorectal cancer. *World J Surg* 2000; 24: 1052-5.

Nomori H, Watanabe K, Ohtsuka T, Naruke T, Suemasu K, Uno K. Evaluation of F-18 fluorodeoxyglucose (FDG) PET scanning for pulmonary nodules less than 3 cm in diameter, with special reference to the CT images. *Lung Cancer* 2004; 45: 19-27.

Nordic Gastrointestinal Tumor Adjuvant Therapy Group. Expectancy of primary chemotherapy in patients with advanced asymptomatic colorectal cancer: a randomized trial. *J Clin Oncol* 1992; 10: 904-911.

Nordlinger B, Sorbye H, Colette L, Glimelius B, Poston GJ, Schlag PM, Rougier P et al. Perioperative chemotherapy with FOLFOX4 and surgery versus surgery alone for resectable liver metastases from colorectal cancer (EORTC Intergroup trial 40983): a randomized controlled trial. *Lancet* 2008; 371: 1007-1016.

Northover J. Follow-up after colorectal cancer resection: Is it worthwhile? In: *Challenges in Colorectal Cancer*. ed. Scholefield JH. 63-79. 2000. Blackwell Science Ltd.

Numminen K, Isoniemi H, Halavaara J et al. Preoperative assessment of focal liver lesions: multidetector computed tomography challenges magnetic resonance imaging. *Acta Radiol* 2005; 46: 9-15.

O'Connell MJ, Laurie JA, Kahn M, Fitzgibbons RJ Jr., Erlichman C, Shepherd L et al. "Prospectively randomized trial of postoperative adjuvant chemotherapy in patients with high-risk colon cancer". *J Clin Oncol*, Vol. 16, No. 1, 1998, pp. 295-300.

O'Connell MJ, Maillbrd JA, Kahn MJ, Macdonald JS, Haller DG, Mayer RJ et al. "Controlled trial of fluorouracil and low-dose leucovorin given for 6 months as postoperative adjuvant therapy for colon cancer". *J Clin Oncol*, Vol. 15, No. 1, 1997, pp. 246-250.

Ohlsson B, Breland U, Ekberg H, Graffner H, Tranberg KG. Follow-up after curative surgery for colorectal carcinoma. Randomized comparison with no follow-up. *Dis Colon Rectum* 1995; 38: 619-626.

Olsen KR, Bojesen SE, Gerdes AM, Lindorff-Larsen K, Bernstein IT. Cost-effectiveness of surveillance programs for families at high and moderate risk of hereditary non-polyposis colorectal cancer. *Int J Technol Assess Health Care* 2007; 23: 89-95.

Ong KE, Leen E. Radiological staging of colorectal liver metastases. *Surg Oncol* 2007; 16: 7-14.

Oster G, Tuden RL, Colditz GA. Prevention of venous thromboembolism after general surgery. Cost-effectiveness analysis of alternative approaches to prophylaxis. *Am J Med* 1987; 82: 889-899.

Osti MF, Padovan FS, Pirolli C, Sbarbati S, Tombolini V, Meli C et al. Comparison between transrectal ultrasonography and computed tomography with rectal inflation of gas in preoperative staging of lower rectal cancer. *Eur Radiol* 1997; 7: 26-30.

Pachler J, Wille-Jørgensen P. Quality of life after rectal resection for cancer, with or without permanent colostomy. *Cochrane Data base Syst Rev* 2004; 3: CD004323.

Painter J, Saunders DB, Bell GD, Williams CB, Pitt R, Bladen J. Depth of insertion at flexible sigmoidoscopy: implications for colorectal cancer screening and instrument design. *Endoscopy* 1999; 31: 227-31.

Pakkastie TE, Ovaska JT, Pekkala ES, Luukkonen PE, Järvinen HJ. A randomised study of colostomies in low colorectal anastomoses. *Eur J Surg* 1997; 163: 929-33.

Papagrigoriadis S, Koreli A. The needs of general practitioners in the follow-up of patients with colorectal cancer. *EJSO* 2001; 27: 541-544.

Park J et al. Radiofrequency Ablation for Metachronous Liver Metastasis from Colorectal Cancer after Curative Surgery. *Ann Surg Oncol* 2008; 15: 227-32.

Park J, MacRae H, Musselman LJ, Rossos P, Hamstra SJ, Wolman S et al. Randomized controlled trial of virtual reality simulator training: transfer to live patients. *Am J Surg* 2007; 194: 205-11.

Patel SC, Tovee EB, Langer B. Twenty-five years experience with radical surgical treatment of carcinoma of the extraperitoneal rectum. *Surgery* 1977; 82: 460-5.

Peeters KC, Marijnen CA, Nagtegaal ID, Kranenbarg EK, Putter H, Wiggers T et al. The TME trial after a median follow-up of 6 years: increased local control but no survival benefit in irradiated patients with respectable rectal carcinoma. *Ann Surg* 2007; 246: 693-701.

Peeters KC, Tollenaar RA, Marijnen CA, Klein Kranenbarg E, Steup WH, Wiggers T et al. Risk factors for anastomotic failure after total mesorectal excision of rectal cancer. *Br J Surg* 2005; 92: 211-6.

Pena-Soria MJ, Mayol JM, nula-Fernandez R, rbeo-Escolar A, Fernandez-Represa JA (2007). Mechanical bowel preparation for elective colorectal surgery with primary intraperitoneal anastomosis by a single surgeon: interim analysis of a prospective single-blinded randomized trial. *J Gastrointest Surg* 11: 562-567.

Persson AV, Davis RJ, Villavicencio JL. Deep venous thrombosis and pulmonary embolism. *Surg Clin North Am* 1991; 71: 1195-1209.

Peschaud F, Cuenod CA, Benoist S, Julié C, Beauchet A, Siauve N et al. Accuracy of magnetic resonance imaging in rectal cancer depends on location of the tumor. *Dis Colon Rectum* 2005; 48: 1603-9.

Petersen VS, Baxter KJ, Love SB et al. Identification of objective pathological prognostic determinants and models of prognosis in Dukes B colon cancer. *Gut* 2002; 51: 65-69.

Petrowsky H, Demartines N, Rousson V, Clavien PA. Evidence-based value of prophylactic drainage in gastrointestinal surgery: a systematic review and meta-analyses. *Ann Surg* 2004; 240: 1074-84.

Peuchot M, Libshitz HI. Pulmonary metastatic disease: radiologic-surgical correlation. *Radiology* 1987; 164: 719-722.

Pfeiffer P, Nielsen D, Bjerregaard J, Qvortrup C, Yilmaz M, Jensen B. Biweekly cetuximab and irinotecan as third-line therapy in patients with advanced colorectal cancer after failure to irinotecan, oxaliplatin and 5-fluorouracil. *Ann Oncol* 2008 June; 19 (6): 1141-5.

Pfeiffer P, Nielsen D, Yilmaz M, iversen A, Vejlo C, Jensen BV. Cetuximab and irinotecan as third line therapy in patients with advanced colorectal cancer after failure of irinotecan, oxaliplatin and 5-fluorouracil. *Acta Oncol* 2007; 46 (5): 697-701.

Pfister DG, Benson AB III, Somerfield MR. Clinical practice. Surveillance strategies after curative treatment of colorectal cancer. *N Engl J Med* 2004; 350: 2375-2382.

Pheby DHF, Levine DF, Pitcher RW, Sheperd NA. Lymph node harvests directly influence the staging of colorectal cancer: evidence from a regional audit. *J Clin Pathol* 2004; 57: 43-7.

Phillips RKS, Hittinger R, Blesowsky L, Fry JS, Fielding LP. Local recurrence following curative surgery for large bowel cancer: I. The overall picture. *Br J Surg* 1984; 71: 12-6.

Phillips RKS, Hittinger R, Fry JS, Fielding LP. Malignant large bowel obstruction. *Br J Surg* 1985; 72: 296-302.

Pietra N, Sarli L, Costi R, Ouchemi C, Grattarola M, Peracchia A. Role of follow-up in management of local recurrences of colorectal cancer: a prospective, randomized study. *Dis Colon Rectum* 1998a; 41: 1127-1133.

Pietra N, Sarli L, Thenasseril BJ, Costi R, Sansebastiano G, Peracchia A. Risk factors of local recurrence of colorectal cancer: a multivariate study. *Hepatogastroenterology* 1998b; 45: 1573-1578.

Pignone M, Rich M, Teutsch SM, Berg AO, Lohr KN. Screening for colorectal cancer in adults at average risk: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2002; 137: 132-41.

Pikarsky A, Wexner S, Lebensart P, Efron J, Weiss E, Noguera J et al. The use of rectal ultrasound for the correct diagnosis and treatment of rectal villous tumors. *Am J Surg* 2000; 179: 261-5.

Piscaglia F, Corradi F, Mancini M et al. Real time contrast enhanced ultrasonography in detection of liver metastases from gastrointestinal cancer. *BMC Cancer* 2007; 7: 171.
Pittam 1984.

Place: The relationship between concentration, patient accessibility and utilization of services. I: Ferguson B, Posnett JW, Sheldon TA, red. Concentration and choice of provision of hospital services. York: NHS Centre for Reviews and Dissemination, University of York, 1997.

Platell C, Barwood N, Makin G (2006). Randomized clinical trial of bowel preparation with a single phosphate enema or polyethylene glycol before elective colorectal surgery. *Br J Surg* 93: 427-433.

Pollack J et al. Late adverse effects of short-course preoperative radiotherapy in rectal cancer. *Br J Surg* 2006; 93: 1519-25.

Pollett WG, Nicholls RJ. The relationship between the extent of distal clearance and survival and local recurrence rates after curative anterior resection for carcinoma of the rectum. *Ann Surg* 1983; 198: 159-63.

Popat S, Hubner R, Houlston RS. Systematic review of microsatellite instability and colorectal cancer prognosis. *J Clin Oncol* 2005; 23 (3): 609-618.

Popin EA, Benedetti JK, Estes NC, Haller DG, Mayer RJ, Goldberg RM et al. "Phase III Southwest Oncology Group 9415/Intergroup 0153 randomized trial of fluorouracil, leucovorin, and levamisole versus fluorouracil continuous infusion and levamisole for

adjuvant treatment of stage III and high-risk stage II colon cancer 1". *J Clin Oncol*, Vol. 23, No. 9, 2005, pp. 1819-1825.

Portier G, Ghouti L, Kirzin S, Guimbaud R, Rives M, Lazorthes F. Oncological outcome of ultra-low coloanal anastomosis with and without intersphincteric resection for low rectal adenocarcinoma. *Br J Surg* 2007; 94: 341-5.

Povoski SP, Fong Y, Sgouros SC, Kemeny NE, Downey RJ, Blumgart LH. Role of chest CT in patients with negative chest x-rays referred for hepatic colorectal metastases. *Ann Surg Oncol* 1998; 5: 9-15.

Prall F et al. Tumour budding as prognostic factor in stage I/II colorectal carcinoma. *Histopathology* 2005; 47: 17-24.

Punt CJ, Tol J, Rodenburg CJ, Cats A, Creemers G, Schrama JG et al. Randomized phase III study of capecitabine, oxliplatin, and bevacizumab with or without cetuximab in advanced colorectal cancer (ACC), the CAIRO2 study of the Dutch Colorectal Cancer Group (DCCG). *proc Am Soc Clin Oncol* 2008; #LBA4011.

Purkayastha S, Tekkis PP, Athanasiou T, Aziz O, Negus R, Gedroyc W et al. Magnetic resonance colonography versus colonoscopy as a diagnostic investigation for colorectal cancer: a meta-analysis. *Clin Radiol* 2005; 60: 980-9.

Quaia E, D'Onofrio M, Palumbo A, Rossi S, Bruni S, Cova M. Comparison of contrast-enhanced ultrasonography versus baseline ultrasound and contrast-enhanced computed tomography in metastatic disease of the liver: diagnostic performance and confidence. *Eur Radiol* 2006; 16: 1599-1609.

Quasar Collaborative Group, Gray R, Barnwell J, McConkey C, Hills RK, Williams NS et al. "Adjuvant chemotherapy versus observation in patients with colorectal cancer: a randomised study". *Lancet*, Vol. 370, No. 9604, 2007, pp. 2020-2029.

Quirke P, Dixon MF. The prediction of local recurrence in rectal adenocarcinoma by histopathological examination. *Int J Colorectal Dis* 1988; 3: 127-31.

Quirke P. Training and quality assurance for rectal cancer: 20 years of data is enough. *Lancet Oncol* 2003; 4: 695-702.

Radcliffe A. Can the results of anorectal (abdominoperineal) resection be improved: are circumferential resection margins too often positive? *Colorectal Dis* 2006; 8: 160-7.

Rafaelsen SR, Kronborg O, Fenger C. Echo pattern of lymph nodes in colorectal cancer. *Br J Radiol* 1992; 62: 218.

Rafaelsen SR, Kronborg O, Larsen C, Fenger C. Intraoperative ultrasonography in detection of hepatic metastases from colorectal cancer. *Dis Colon Rectum* 1995; 38: 355-360.

Rafaelsen SR, Kronborg O, Larsen C, Fenger C. Intraoperative ultrasonography in detection of hepatic metastases from colorectal cancer. *The Year Book of Surgery* 1996a; 11: 412-5.

Rafaelsen SR, Sørensen T, Jakobsen A, Bisgaard C, Lindebjerg J. Transrectal ultrasonography and magnetic resonance imaging in the staging of rectal cancer. Effect of experience. *Scand J Gastroenterol* 2008; 43: 400-6.

Ram E, Sherman Y, Weil R, Vishne T, Kravarusic D, Dreznik Z (2005). Is mechanical bowel preparation mandatory for elective colon surgery? A prospective randomized study. *Arch Surg* 140: 285-288.

Ramakrishnan K, Scheid DC. Selecting patients for flexible sigmoidoscopy. Determinants of incomplete depth of insertion. *Cancer* 2005; 103: 1179-85.

Rappeport ED, Loft A, Berthelsen AK et al. Contrast-enhanced FDG-PET/CT vs. SPIO-enhanced MRI vs FDG-PET vs CT in patients with liver metastases from colorectal cancer: a prospective study with intraoperative confirmation. *Acta Radiol* 2007; 48: 369-378.

Rasmussen MS, Jørgensen LN, Wille-Jørgensen P, Nielsen JD, Horn A, Mohn-A-C et al. Prolonged prophylaxis with low dalteparin to prevent late thromboembolic complications in patients undergoing major abdominal surgery. A multicenter randomized open-label study. *J Thromb Haemostas* 2006; 4: 2384-2390.

Rasmussen PC. Primary Advanced and Recurrent Rectal Cancer. *Dansk Kirurgisk Selskabs 100 års Jubilæumsmøde*, 2008.

Ratto C, Sofo L, Ippoliti M et al. Accurate lymph-node detection in colorectal specimens resected for cancer is of prognostic significance. *Dis Colon Rectum* 1999; 42 (2): 143-54.

Raue W, Haas O, Junghans T, Scharfenberg M, Müller J, Schwenk W (2004). "Fast-track" multimodal rehabilitation program improves outcome after laparoscopic sigmoidectomy: a controlled prospective evaluation. *Surg Endosc* 18: 1463-1468.

Rembacken B. Missing colorectal cancer. *Endoscopy* 2004; 36: 551-553.

Rehnan AG, Egger M, Saunders MP, O'Dwyer ST. Impact on survival of intensive follow-up after curative resection for colorectal cancer: systematic review and meta-analysis of randomised trials. *BMJ* 2002; 324: 813.

Rehnan AG, O'Dwyer ST, Whynes DK. Cost effectiveness analysis of intensive versus conventional follow-up after curative resection for colorectal cancer. *BMJ* 2004; 328: 81.

Renkonen-Sinisalo L, Kivisaari A, Kivisaari L, Sarna S, Järvinen HJ. Utility of computed tomographic colonography in surveillance for hereditary nonpolyposis colorectal cancer syndrome. *Fam Cancer* 2007; 6: 135-140.

Reuter NP et al. Radiofrequency Ablation vs. Resection for Hepatic Colorectal Metastasis: Therapeutically Equivalent? *J Gastrointest Surg* 2008 Oct 30. (Epub ahead of print).

Reza MM, Blasco JA, Andradas E, Cantero R, Mayol J. Systematic review of laparoscopic versus open surgery for colorectal cancer. *Br J Surg* 2006; 93: 921-928.

Ribic DM, Sargent DJ, Moore MJ et al. Tumor microsatellite-instability status as a predictor of benefit from fluorouracil-based adjuvant chemotherapy for colon cancer. *N Engl J Med* 2003; 349: 247-57.

Richards CJ, West KP. Rapid turnaround in histopathology is not appropriate for colorectal resections. *J Pathol* 1998; 186: (suppl) 29A.

Rifkin MD, Ehrlich SM, Marks G. Staging of rectal carcinoma: Prospective comparison of endorectal US and CT. *Radiology* 1989; 170: 319.

Robinson MH, Marks CG, Farrands PA, Thomas WM, Hardcastle JD. Population screening for colorectal cancer: comparison between guaiac and immunological faecal occult blood tests. *Br J Surg* 1994; 81: 448-51.

Rodriguez-Moranta F, Salo J, Arcusa A et al. Postoperative surveillance in patients with colorectal cancer, who have undergone curative resection: a prospective multicenter, randomized, controlled trial. *J Clin Oncol* 2006; 24: 386-393.

Rosa F, Meimarakis G, Stahl A et al. Colorectal cancer patients before resection of hepatic metastases. Impact of (18)F-FDG PET on detecting extrahepatic disease. *Nuklearmedizin* 2004; 43: 135-140.

Rosman AS, Kortsen MA. Meta-analysis comparing CT colonography, air contrast barium enema, and colonoscopy. *Am J Med* 2007 Mar; 120 (3): 203-210.

Rothenberg ML, Oza AM, Bigelow RH, Berlin JD, Marshall JL, Ramanathan RK et al. Superiority of exaliplatin and fluorouracil-leucovorin compared with either therapy alone in patients with progressive colorectal cancer after irinotecan and fluorouracil-leucovorin: interim results of a phase III trial. *J Clin Oncol* 2003 June 1; 21 (11): 2059-69.

Rouffet F, Hay JM, Vacher B, Fingerhut A, Elhadad A, Flamant Y et al. Curative resection for left colonic carcinoma: Hemicolectomy vs. segmental colectomy. A prospective, controlled, multicenter trial. *French Association for Surgical Research. Dis Colon Rectum* 1994; 37: 651-9.

Rovera F, Dionigi G, Boni L, Piscopo C, Masciocchi P, Alberio MG et al. 2007. Infectious complications in colorectal surgery. *Surg Oncol* 16 Suppl 1: S121-S124.

Rowland M, Tozer T. 1989. *Clinical Pharmacokinetics*. Second Edition p. 78-100. In Lea & Fibiger (ed.), Pennsylvania.

Rubio CA, Jaramillo E, Lindblom A, Fogt F. Classification of colorectal polyps: Guidelines for the Endoscopist. *Endoscopy* 2002; 34: 226-36.

Ruers TJ, Langenhoff BS, Neeleman N et al. Value of positron emission tomography with (F-18) fluorodeoxyglucose in patients with colorectal liver metastases: a prospective study. *J Clin Oncol* 2002; 20: 388-395.

Rullier E, Laurent C, Carles J, Saric J, Michel P, Parneix M. Local recurrence of low rectal cancer after abdominoperineal and anterior resection. *Br J Surg* 1997; 84: 525-8.

Rullier E, Laurent C, Garrelon JL, Michel P, Saric J, Parneix M. Risk factors for anastomotic leakage after resection of rectal cancer. *Br J Surg* 1998; 85: 355-8.

Rullier E, Laurent C, Bretagnol F, Rullier A, Vendrely V, Zerbib F. Sphincter-saving resection for all rectal carcinomas: the end of the 2-cm distal rule. *Ann Surg* 2005; 241: 465-9.

Runkel NS, Schlag P, Schwarz V, Herfarth C. Outcome after emergency surgery for cancer of the large intestine. *Br J Surg* 1991; 78: 183-8.

Ruo L, Gougoutas C, Paty PB, Guillem JG, Cohen AM, Wong WD. *J Am Coll Surg* 2003; 196: 722-8.

Saclarides TJ, Bhattacharyya AK, Britton-Kuzel C, Szeluga D, Economou SG. Predicting lymph node metastases in rectal cancer. *Dis Colon Rectum* 1994; 37: 52-7.

Saha S, Dan AG, Beutler T, Wiese D, Schochet E, Badin J et al. Sentinal lymph node mapping technique in colon cancer. *Semin Oncol* 2004; 31: 374-81.

Saida Y, Sumiyama Y, Nagao J et al. Long-term prognosis of preoperative "bridge to surgery" expandable metallic stent insertion for obstructive colorectal cancer. Comparison with emergency operation. *Dis Colon Rectum* 2003; 46 (suppl): S44-S49.

Sailer M, Fuchs KH, Fein M, Thiede A. Randomized clinical trial comparing quality of life after straight and pouch coloanal reconstruction. *Br J Surg* 2002; 89: 1108-17.

Sakamoto J, Hamada C, Yoshida S et al. An individual patient data meta-analysis of adjuvant therapy with uracil-tegafur (UFT) in patients with curatively resected rectal cancer. *Br J Cancer* 2007; 96: 1170-1177.

Sales J-P, Wind P, Douard R, Cugnenc P-H, Loric S. No-touch technique is used to minimise spread of malignant cells during surgery for colorectal cancer. *Lancet* 1999; 354-92.

Salomon S, Jensen TG, Qvist N, Frimodt-Moller N, Pedersen C, Madsen H. 2007. Use of antibiotics in colorectal surgery in Denmark. *Ugeskr Laeger* 169: 920-925.

Salovaara R, Loukola A, Kristo P, Kaariainen H, Ahtola H, Eskelinen M et al. Population-based molecular detection of hereditary nonpolyposis colorectal cancer. *J Clin Oncol* 2000; 18: 2193-2200.

Saltz LB, Clarke S, az-Rubio E, Scheithauer W, Figer A, Wong R et al. Bevacizumab in combination with oxaliplatin-based chemotherapy as first-line therapy in metastatic colorectal cancer: a randomized phase III study. *J Clin Oncol* 2008 April 20; 26 (12): 2013-9.

Saltz LB, Cox JV, Blanke C, Rosen LS, Fehrenbacher L, Moore MJ et al. Irinotecan plus fluorouracil and leucovorin for metastatic colorectal cancer. Irinotecan Study Group. *N Engl J Med* 2000 September 28; 343 (13): 905-14.

Saltz LB. Irinotecan in the first-line treatment of colorectal cancer. *Oncology (williston Park)* 1998 August; 12 (8 Suppl 6): 54-8.

Santos JCM Jr., Batista J, Sirimarco MT, Guimaraes AS, Levy CE (1994). Prospective randomized trial of mechanical bowel preparation in patients undergoing elective colorectal surgery. *Br J Surg* 81: 1673-1676.

Sargent DJ, Goldberg RM, Jacobson SD, Macdonald JS, Lablbnca R, Haller DG et al. "A pooled analysis of adjuvant chemotherapy for resected colon cancer in elderly patients 1". *N Engl J Med*, Vol. 345, No. 15, 2001, pp. 1091-1097.

Sato K, Inomata M, Kakisako K, Shiraishi N, Adachi Y, Kitano S. Surgical technique influences bowel function after low anterior resection and sigmoid colectomy. *Hepatogastroenterology* 2003; 50: 1381-4.

Sayfan J, Averbuch F, Koltun L, Benyamin N. Effect of rectal stump washout on the presence of free malignant fcells in the rectum during anterior resection for rectal cancer. *Dis Colon Rectum* 2000; 43: 1710-2.

Schaefer O, Langer M. Detection of recurrent rectal cancer with CT, MRI and PET/CT. *Eur Radiol* 2007; 17: 2044-54.

Scheithauer W, McKendrick J, Begbie S, Borner M, Burns WI, Burris HA et al. "Oral capecitabine as an alternative to iv 5-fluorouracil-based adjuvant therapy for colon cancer: safety results of a randomized, phase III trial". *Ann Oncol*, Vol. 14. No. 12, 2003, pp. 1735-1743.

Scheithauer W, Rosen H, Kornek GV, Sebesta C, Depisch D. Randomised comparison of combination chemotherapy plus supportive care with supportive care alone in patients with metastatic colorectal cancer. *BMJ* 1993; 306: 752-755.

Schiesser M, Muller S, Kirchhoff P, Breintenstein S, Schafer M, Clavien PA (2008). Assessment of a novel screening score for nutritional risk in predicting complications in gastro-intestinal surgery. *Clin Nutr* 2008; 27: 565-70.

Schilder JM, Hurteau JA, Look KY, Moore DH, Raff G, Stehman FB et al (1997). A prospective controlled trial of early postoperative oral intake following major abdominal gynecologic surgery. *Gynecol Oncol* 67: 235-240.

Schlachta CM, Mamazza J, Seshadri PA, Cadeddu M, Gregoire R, Poulin EC. Defining a learning curve for laparoscopic colorectal resections. *Dis Colon Rectum* 2001; 44: 217-222.

Schmidt O, Merkel S, Hohenberger W. Anastomotic leakage after low rectal stapler anastomosis: Significance of intraoperative anastomotic testing. *Eur J Surg Oncol* 2003; 29: 239-43.

Schmidt J, Strotzer M, Fraunhofer S, Boedeker H, Zirngibl H. Intraoperative ultrasonography versus helical computed tomography and computed tomography with arteriography in diagnosing colorectal liver metastases: lesion-by-lesion analysis. *World J Surg* 2000; 24: 43-47; discussion 48.

Schmoll HJ, Cartwright T, Tabernero J, Nowacki MP, Figer A, Maroun J et al. "Phase III trial of capecitabine plus oxaliplatin as adjuvant therapy for stage III colon cancer: a planned safety analysis in 1,864 patients 1". *J Clin Oncol*, Vol. 25, No. 1, 2007, pp. 102-109.

Schoemaker D, Black R, Giles L, Toouli J. Yearly colonoscopy, liver CT, and chest radiography do not influence 5-year survival of colorectal cancer patients. *Gastroenterology* 1998; 114: 7-14.

Schulze S, Lyng K-K. Palliation of rectosigmoid neoplasms with Nd: YAG laser treatment. *Dis Colon Rectum* 1994; 37: 882-4.

Schwenk W, Haase O, Neudecker J, Muller JM. Short term benefits for laparoscopic colorectal resection. *Cochrane Database Syst Rev* 2005 (3): CD003145.

Schäfer H, Baldus SE, Casper F, Hölscher AH. Submucosal infiltration and local recurrence in pT1 low-risk rectal cancer treated by transanal endoscopic microsurgery. *Chirurg* 2005; 76: 379-84.

Screening for tarmkræft i Vejle amt – en gennemførlighedsundersøgelse august 2005 – november 2006. 2007.

Secco GB, Fardelli R, Gianquinto D, Bonfante P, Baldi E, Ravera G et al. Efficacy and cost of risk-adapted follow-up in patients after colorectal cancer surgery: a prospective, randomized and controlled trial. *Eur J Surg Oncol* 2002; 28: 418-423.

Sekimoto M. Laparoscopic resection for colorectal cancer in Japan. *Dis Colon Rectum* 2007; 50: 1708-1714.

Seow-Choen F, Goh HS. Prospective randomized trial comparing J colonic pouch-anal anastomosis and straight coloanal reconstruction. *Br J Surg* 1995; 82: 608-10.

Shah HA, Paszat LF, Saskin R, Stukel TA, Rabeneck L. Factors associated with incomplete colonoscopy: a population-based study. *Gastroenterology* 2007; 132: 2297-303.

Shibata D, Guillem JG, Lanouette N, Paty P, Minsky B, Harrison L et al. Functional and quality-of-life outcomes in patients with rectal cancer after combined modality therapy, intraoperative radiation therapy, and sphincter preservation. *Dis Colon Rectum* 2000; 43: 752-8.

Shihab OC, Moran BJ, Heald RJ, Quirke P, Brown G. MRI staging of low rectal cancer. *Eur Radiol* 2008.

Shinto E et al. A novel classification of tumour budding in colorectal cancer based on the presence of cytoplasmic pseudo-fragments around budding foci. *Histopathology* 2005; 47: 25-31.

Sigounas D, Christodoulou D, Marcon N, Tsianos EV. Endoscopic Mucosal Resection – A Comprehensive Review. *Gastric Breast Cancer* 2004; 3 (1): 27-35.

Simmonds PC. Palliative chemotherapy for advanced colorectal cancer: systematic review and meta-analysis. Colorectal Cancer Collaborative Group. *BMJ* 2000; 321: 531-535.

Sisisriro R, Boonkitticharoen V, Kraiphibul P, Ratanatharathorn V, Sumboonnanon K, Kanjanapitak A et al. Detection of colorectal carcinoma by anti-CEA monoclonal antibody (IOR-CEA1) labeled with ^{99m}Tc scintigraphy. *Hepatogastroenterology* 2000; 47: 405-413.

Sjödahl R, Franzen T, Nyström PO. Primary versus staged resection for acute obstructing colorectal carcinoma. *Br J Surg* 1992; 79: 685-8.

Skandarajah AR, Tjandra JJ. Preoperative loco-regional imaging in rectal cancer. *ANZ J Surg* 2006; 76: 497-504.

Skånberg J, Lundholm K, Haglind E. Effects of blood transfusion with leucocyte depletion on length of hospital stay, respiratory assistance and survival after curative surgery for colorectal cancer. *Acta Oncol* 2007; 46 (8): 1123-30.

Slim K, Vicaut E, Panis Y, Chipponi J (2004). Meta-analysis of randomized clinical trials of colorectal surgery with or without mechanical bowel preparation. *Br J Surg* 91: 1125-1130.

Smedh K, Olsson L, Johansson H et al. Reduction of postoperative morbidity and mortality in patients with rectal cancer following the introduction of a colorectal unit. *Br J Surg* 2001; 88: 273-7.

Smedley F, Bowling T, James M, Stokes E, Goodger C, O'Connor O et al (2004). Randomized clinical trial of the effects of preoperative and postoperative oral nutritional supplements on clinical course and cost of care. *Br J Surg* 91: 983-990.

Smith EH. Complications of percutaneous abdominal fine-needle biopsy. *Radiology* 1991; 178: 253-258.

Sobin LH, Wittekind C and International Union Against Cancer (UICC). TNM classification on malignant tumours (5th edn.), Wiley-Liss Publications, New York (1997).

Sobin LH, Wittekind C. UICC TNM Classification on malignant tumors, 6 th edn. New York: Wiley-Liss 2002.

Sobrero AF, Maurel J, Fehrenbacher L, Scheithauer W, Abubakr YA, Lutz MP et al. EPIC: phase III trial of cetuximab plus irinotecan after fluoropyrimidine and oxaliplatin failure in patients with metastatic colorectal cancer. *J Clin Oncol* 2008 May 10; 26 (14): 2311-9.

Society of American Gastrointestinal and Endoscopic Surgeons (SAGES). Guidelines for laparoscopic resection of curable colon and rectal cancer. SAGES 2006.

Song F, Glenny AM. 1998. Antimicrobial prophylaxis in colorectal surgery: a systematic review of randomised controlled trials. *Health Techno Assess* 2: 1-100.

Soop M, Nygren J, Myrenfors P, Thorell A, Ljungqvist O (2001). Preoperative oral carbohydrate treatment attenuated immediate postoperative insulin resistance. *Am J Physiol Endocrinol Metab* 280: E576-E583.

Sosna J, Sella T, Sy O, Lavin PT, Eliahou R, Fraifeld S et al. Critical analysis of the performance of double-contrast barium enema for detecting colorectal polyps \geq 6 mm in the era of CT colonography. *Am J Roentgenol* 2008; 190: 374-85.

Sowden AJ, Grilli R, Rice N. The relationship between hospital volume and quality of health outcomes. I: Ferguson B, Posnett JW, Sheldon TA, red. Concentration and choice of provision of hospital services. York: NHS Centre for Reviews and Dissemination, University of Yourk, 1997.

Sprangens M, Te Velde A, Aaronson K, Taal B. Quality of life following surgery for colorectal cancer. *Psycho-Oncology* 1993; 2: 247-259.

Starck M, Bohe M, Fork FT et al. Endoluminal ultrasound and low-field magnetic resonance imaging are superior to clinical examination the preoperative staging of rectal cancer. *Eur J Surg* 1995; 161: 841-5.

Statens Institut for Medicinsk Teknologivurdering. Kræft i tyktarm og endetarm. Diagnostik og screening. Medicinsk Teknologivurdering 2001; 3 (1).

Stelzner M. 2003 SSAT-AGA-ASGE Workshop on "Palliative Therapy of Rectal Cancer". Summary statement. *J Gastrointest Surg* 2004; 8: 253-8.

Stephenson BM, Shandall AA, Farouk R, Griffith G. Malignant left-sided large bowel obstruction managed by subtotal/total colectomy. *Br J Surg* 1990; 77: 1098-102.

Sterk P, Shekarriz B, Günter S, Nolde J, Kehller R, Bruch HP et al. Voiding and sexual dysfunction after deep rectal resection and total mesorectal excision: prospective study on 52 patients. *Int J Colorectal Dis* 2005; 20: 423-7.

Stewart J, Finan PJ, Courtney DF, Brennan TG. Does a water-soluble contrast enema assist in the management of acute large bowel obstruction: A prospective study of 117 cases. *Br J Surg* 1984; 71: 799-801.

Stewart CJ, Morris M, de Boer B, Iacopetta B. Identification of serosal invasion and extramural venous invasion on review of Dukes' stage B colonic carcinomas and correlation with survival. *Histopathol* 2007; 51: 372-8.

Stiggelbout AM, de Haes J, Vree R, van de Velde C, Bruijninx C, van Groningen K et al. Follow-up of colorectal cancer patients: quality of life and attitudes towards follow-up. *Br J Surg* 1997; 75: 914-920.

Stipa F, Lucandri G, Ferri M, Casula G, Ziparo V. Local excision of rectal cancer with transanal endoscopic microsurgery (TEM). *Anticancer Res* 2004; 24: 1167-72.

Stocchi L, Nelson H. Diagnostic and therapeutic applications of monoclonal antibodies in colorectal cancer. *Dis Colon Rectum* 1998; 41: 232-250.

Stoker J, Rociu E, Wiersma TG, Laméris JS. Imaging of anorectal disease. *Br J Surg* 2000; 87: 10-27.

Stratton RJ, Elia M (2007). Who Benefits from nutritional support: what is the evidence? *Eur J Gastroenterol Hepatol* 19: 353-358.

Sundhedsstyrelsen. Behandling af tarmkræft i Danmark med fokus på den kirurgiske behandling. www.sst.dk. Sundhedsstyrelsen 2004.

Sundhedsstyrelsen, Monitorering & Medicinsk Teknologivurdering. Screening for tarmkræft: Deltagelsesprocentens betydning – En medicinsk teknologivurdering. København: Sundhedsstyrelsen, Medicinsk Teknologivurdering 2008; 10 (1).

Sungurtekin H, Sungurtekin U, Balci C, Zencir M, Erdem E (2004). The influence of nutritional status on complications after major intraabdominal surgery. *J Am Coll Nutr* 23: 227-232.

Svendsen LB, Bülow S, Mellempgaard A. Metachronous colorectal cancer in young patients: Expression of the hereditary nonpolyposis colorectal cancer syndrome? *Dis Colon Rectum* 1991; 34: 790-793.

Svensson MH, Svensson E, Lasson A, Helström M. Patient acceptance of CT colonography and conventional colonoscopy: Prospective comparative study in patients with or suspected of having colorectal disease. *Radiology* 2002; 222: 337-45.

Swedish Rectal Cancer Trial. Improved Survival With Preoperative Radiotherapy in Resectable Rectal Cancer. *N Engl J Med* 1997; 336: 980-987.

Swensen SJ, Jett JR, Hartman TE et al. CT screening for lung cancer: five-year prospective experience. *Radiology* 2005; 235: 259-265.

Sørensen LT, Jørgensen T, Kirkeby LT et al. Smoking and alcohol abuse are major risk factors for anastomotic leakage in colorectal surgery. *Br J Surg* 1999; 86: 927-31.

Takeuchi N, Ramirez JM, Mortensen NJ, Cobb R, Whittlestone T. Intraoperative ultrasonography in the diagnosis of hepatic metastases during surgery for colorectal cancer. *Int J Colorectal Dis* 1996; 11: 92-95.

Tang R, Chen HH, Wang YL, Changchien CR, Chen JS, Hsu KC et al. Risk factors for surgical site infection after elective resection of the colon and rectum: a single-center prospective study of 2,809 consecutive patients. *Ann Surg* 2001 Aug; 234 (2): 181-9.

Tarantino I, Hetzer FH, Warschkow R, Zünd M, Stein HJ, Zerz A. Local excision and endoscopic posterior mesorectal resection versus low anterior resection in T1 rectal cancer. *Br J Surg* 2008; 95: 375-80.

Taylor SA, Halligan S, Saunders BP, Bassett P, Vance M, Bartram CI. Acceptance by patients of multidetector CT colonography compared with barium enema examinations, flexible sigmoidoscopy, and colonoscopy. *Am J Roentgenol* 2003; 181: 913-21.

Taylor SA, Halligan S, Burling D, Roddie ME, Honeyfield L, McQuillan J et al. Computer-assisted reader software versus expert reviewers for polyp detection on CT colonography. *Am J Roentgenol* 2006; 186: 696-702.

Taylor SA, Suzuki N, Beddoe G, halligan S. Flat neoplasia of the colon: CT colonography with CAD. *Abdom Imaging* 2008 May 30.

Tei TM, Stolzenburg T, Buntzen S, Laurberg S, Kjeldsen H. Use of transpelvic rectus abdominis musculocutaneous flap for anal cancer salvage surgery. *Br J Surg* 2003 May; 90 (5): 575-80.

Tekkis PP, Senagore AJ, Delaney CP, Fazio VW. Evaluation of the learning curve in laparoscopic colorectal surgery: comparison of right-sided and left-sided resections. *Ann Surg* 2005; 242: 83-91.

Terzi C, Unek T, Sagöl O, Yilmaz T, Füzün M, Sökmen S et al. Is rectal washout necessary in anterior resection for rectal cancer? A prospective clinical study. *World J Surg* 2006; 30: 233-41.

The Clinical Outcomes of Surgical Therapy Study Group. A comparison of laparoscopically assisted and open colectomy for colon cancer. *N Engl J Med* 2004; 350: 2050-2059.

The SCOTIA Study Group. Single-stage treatment for malignant left-sided colonic obstruction: a prospective randomized clinical trial comparing subtotal colectomy with segmental resection following intraoperative irrigation. *Subtotal Colectomy versus On-table Irrigation and Anastomosis*.

Thiis-Evensen E, Hoff GS, Sauar J, Langmark F, Majak BM, Vatn MH. Population-based surveillance by colonoscopy: effect on the incidence of colorectal cancer. Telemark Polyp Study I. *Scand J Gastroenterol* 1999; 34: 414-20.

Thirion P, Michiels S, Pignon JP, Buyse M, Braud AC, Carlson RW et al. Modulation of fluorouracil by leucovorin in patients with advanced colorectal cancer: an updated meta-analysis. *J Clin Oncol* 2004; 22: 3766-75.

Thompson MR, Perera R, Senapati A, Dodds S. Predictive value of common symptom combinations in diagnosing colorectal cancer. *Br J Surg* 2007; 94: 1260-5.

Thomson WHF, Foy CJW, Longman RJ. The nature of local recurrence after colorectal cancer resection. *Colorectal Dis* 2007; 10: 69-71.

Ticani E, piccolo M, Turrini F, Crowther MA, Melotti G, Bondi M. Video laparoscopic surgery: is out-of hospital thromboprophylaxis necessary? *J Thromb Haemost* 2005; 3: 216-20.

Tilney HS, Heriot AG, Purkayastha S, Antoniou A, Aylin P, Darzi AW et al. A national perspective on the decline of abdominoperineal resection for rectal cancer. *Ann Surg* 2008; 247: 77-84.

Tilney HS, Tekkis PP. Extending the horizons of restorative rectal surgery: intersphincteric resection for low rectal cancer. *Colorectal Dis* 2008; 10: 3-15.

Titu LV, Tweedle E, Rooney PS. High tie of the inferior mesenteric artery in curative surgery for left colonic and rectal cancers: a systematic review. *Dig Surg* 2008; 25: 148-57.

Tjandra JJ et al. Follow-Up after Curative Resection of Colorectal Cancer: A Meta-Analysis. *Dis Colon Rectum* 2007; 50: 1783-1799.

Tjandra JJ, Chan MK. Systematic review on the short-term outcome of laparoscopic resection for colon and rectosigmoid cancer. *Colorectal Dis* 2006; 8: 375-388.

Tjandra JJ, Chan MKY. Follow-up after curative resection of colorectal cancer. A meta-analysis. *Dis Colon Rectum* 2007; 50: 1783-1799.

Tjandra JJ, Skandarajak AR. Preoperative loco-regional imaging in rectal cancer. *ANZ J Surg* 2006; 76: 497-504.

Tocchi A, Mazzone G, Lepre L, Costa G, Liotta G, Agostini N et al. Prospective evaluation of omentoplasty in preventing leakage of colorectal anastomosis. *Dis Colon Rectum* 2000; 43: 951-5.

Torngren S, Rieger A. Prophylaxis of deep venous thrombosis in colorectal surgery. *Dis Colon Rectum* 1982; 25: 563-566.

Torricelli P, Pecchi A, Luppi G, Romagnoli R. Gadolinium-enhanced MRI with dynamic evaluation in diagnosing the local recurrence of rectal cancer. *Abdom Imaging* 2003; 28: 19-27.

Tournigand C, Andre T, Achille E, Lledo G, Flesch M, Mery-Mignard D et al. FOLFIRI followed by FOLFOX6 or the reverse sequence in advanced colorectal cancer: a randomized GERCOR study. *J Clin Oncol* 2004 January 15; 22 (2): 229-37.

Tournigand C, Vervantes A, Figer A, Lledo G, Flesch M, Buyse M et al. OPTIMOX1: a randomized study of FOLFOX4 or FOLFIRI with oxaliplatin in a stop-and-go fashion in advanced colorectal cancer – a GERCOR study. *J Clin Oncol* 2006 January 20; 24 (3): 394-400.

Towler B, Irwig L, Glasziou P, Kewenter J, Weller D, Silagy C. A systematic review of the effects of screening for colorectal cancer using the faecal occult blood test, Hemoccult. *BMJ* 1998; 317: 559-65.

Tran DQ, Rosen L, Kim R, Riether RD, Stasik JJ, Khubchandani IT. Actual colonoscopy: What are the risk of perforation? *Am Surg* 2001; 67: 845-8.

Trent/Wales Audit. Guidelines for the Management of Colorectal Cancer, The Royal College of Surgeons of England & Association of Coloproctology of Great Britain and Ireland 1996.

Turnbull RB, Kyle K, Watson FB, Spratt J. Cancer of the colon: The influence of the no-touch isolation technic on survival rates. *Ann Surg* 1967; 166: 420-7.

Tutt AN, Plunkett TA, Barrington SF, Leslie MD. The role of positron emission tomography in the management of colorectal cancer. *Colorectal Dis* 2004; 6: 2-9.

Twelves C, Gollins S, Grieve R, Samuel L. A randomised cross-over trial comparing patient preference for oral capecitabine and 5-fluorouracil/leucovorin regimens in patients with advanced colorectal cancer. *Ann Oncol* 2006; 17: 239-45.

Tytherleigh MG, McC Mortensen NJ. Options for sphincter preservation in surgery for low rectal cancer. *Br J Surg* 2003; 90: 922-33.

Tytherleigh MG, Ng W, Piattathankal AA, Wilson MJ, Farouk R. Preoperative staging of rectal cancer by magnetic resonance imaging remains an imprecise tool. *ANZ J Surg* 2008; 78: 194-8.

Tøttrup A, Frost L. Pelvic sepsis after extended Hartmann's procedure. *Dis Colon Rectum* 2005; 48: 251-5.

Taal BG, van Tinteren H, Zoetmulder FA et al. Adjuvant 5-FU plus levamisol in colonic or rectalcancer: improved survival in stage II and III. *Br J Cancer* 2001; 85: 1437-1443.

Ueno H, Mochizuki H, Hashiguchi Y, Ishikawa K, Fujimoto H, Shinto E et al. Preoperative parameters expanding the indication of sphincter preserving surgery in patients with advanced low rectal cancer. *Ann Surg* 2004; 239: 34-42.

UICC TNM Supplement, 2nd ed (Wittekind Ch, Henson DE, Hutter RVP, Sobin LH) 2001. Wiley-Liss, New York.

UK – Colorectal – Cancer – Screening – Pilot – Group. Results of the first round of a demonstration pilot of screening for colorectal cancer in the United Kingdom. *BMJ* 2004; Jul 17: 329 (458): 133.

UK Flexible Sigmoidoscopy Screening Trial Investigators. Single flexible sigmoidoscopy screening to prevent colorectal cancer: baseline findings of a UK multicentre randomised trial. *Lancet* 2002; 359: 1291-300.

UKCCCR. United Kingdom Coordination Committee on Cancer Research. Handbook for the clinico-pathological assessment and staging of colorectal cancer. UKCCCR 1997.

Umpleby HC, Williamson RC. Survival in acute obstructing colorectal carcinoma. *Dis Colon Rectum* 1984; 27: 299-304.

Urbach DR, Kennedy ED, Cohen MM. Colon and rectal anastomoses do not require routine drainage: a systemic review and metaanalyses. *Ann Surg* 1999; 229: 174-80.

Vamvakas EC. White-blood-cell-containing allogeneic blood transfusion and postoperative infection or mortality: an updated meta-analysis. *Vox Sang* 2007; 92: 224-32.

Van Cutsem E, Peeters M, Siena S, Humblet Y, Hendlisz A, Neyns B et al. Open-label phase III trial of panitumumab plus best supportive care compared with best supportive care alone in patients with chemotherapy-refractory metastatic colorectal cancer. *J Clin Oncol* 2007 May 1; 25 (13): 1658-64.

Van Cutsem E, Twelves C, Cassidy J, Allman D, Bajetta E, Boyer M et al. Oral capecitabine compared with intravenous fluorouracil plus leucovorin in patients with metastatic colorectal cancer: results of a large phase III study. *J Clin Oncol* 2001 November 1; 19 (21): 4097-106.

Van Cutsem EJ, Oliveira J. "Colon cancer: ESMO clinical recommendations for diagnosis, adjuvant treatment and follow-up²". *Ann Oncol* Vol. 19, Suppl 2, 2008, pp. ii29-ii30.

Van Dalen R, Church J, McGannon E, Fay S, Burke C, Clark B. Patterns of surgery in patients belonging to Amsterdam-positive families. *Dis Colon Rectum* 2003; 46: 717-620.

Van Dalen R, Church J, McGannon E, Fay S, Burke C, Clark B. Patterns of surgery in patients belonging to Amsterdam-positive families. *Dis Colon Rectum* 2003; 46: 717-620.

van de Watering LM, Brand A, Houbiers JG, Klein Kranenbarg WM, Hermans J, van de Velde C et al. Perioperative blood transfusions, with or without allogeneic leucocytes, relate to survival, not to cancer recurrence. *Br J Surg* 2001; 88: 267-72.

Vasen HF, Moslein G, Alonso A, Bernstein I, Bertario L, Blanco I et al. Guidelines for the clinical management of Lynch syndrome (hereditary on-polyposis cancer). *J Med Genet* 2007; 44: 353-362.

Vasen HF, Watson P, Mecklin JP, Lynch HT. New clinical criteria for hereditary nonpolyposis colorectal cancer (HNPCC, Lynch syndrome) proposed by the International Collaborative group on HNPCC. *Gastroenterol* 1999; 116: 1453-1456.

Veldkamp R, Kuhry E, Hop WC, Jeekel J, Kazemier , Bojer HJ et al. Laparoscopic surgery versus open surgery for colon cancer: short-term outcomes of a randomised trial. *Lancet Oncol* 2005; 6: 477-484.

Vigder L, Tzur N, Huber M, Mahagna M, Amir I. Management of obstructive carcinoma of the left colon. *Arch Surg* 1985; 120: 825-8.

Vignali A, Fazio VW, Lavery IC et al. Factors associated with the occurrence of leaks in stapled rectal anastomoses: a review of 1,014 patients. *J Am Coll Surg* 1997; 185: 105-13. Viiala 2003.

Vliegen R, Dresen R, Beets G, Daniels-Gooszen A, Kessels A, van Engelshoven J et al. The accuracy of Multi-detector row CT for the assessment of tumour invasion of the mesorectal fascia in primary rectal cancer. *Abdom Imagin* 2008; 33: 604-10.

Von Wagner C, Knight K, Halligan S, Atkin W, Lilford R, Morton D et al. Patient experiences of colonoscopy, barium enema and CT colonography: a qualitative study. *Br J Radiol* 2008.

Votrubova J, Belohlavek O, Jaruskova M, Oliverius M, Lohynska R, Tskova K et al. The role of FDG-PET/CT in the detection of recurrent colorectal cancer. *Eur J Nucl Med Mol Imaging* 2006; 33: 779-984.

Vucelic B, Rex D, Pulanic R, Pfefer J, Hrstic I, Levin B et al. The aero-scope: proof of concept of a pneumatic, skill-independent, self-propelling, self-navigation colonoscope. *Gastroenterology* 2006; 130: 672-7.

Waizer A, Powsner E, Russo I, Hadar S, Cytron S, Lombrozo R. Prospective comparative study of magnetic resonance imaging versus transrectal ultrasound for preoperative staging and follow-up of rectal cancer. Preliminary report. *Dis Colon Rectum* 1991; 34: 1068-1072.

Walker KG, Bell SW, Rickard MJ, Mehanna D, Dent OF, Chapuis PH et al. Anastomotic leakage is predictive of diminished survival after potentially curative resection for colorectal cancer. *Ann Surg* 2004; 240: 255-9.

Wallace MB. Intraluminal endoscopic surgery: the Scioto returns. *Gastroenterology* 2007 Mar; 132 (3): 848-52.

Wang H-S, Lin J-K, Mou C-Y, Lin T-C, Chen W-S, Jiang J-K et al. Long-term prognosis of patients with obstructing carcinoma of the right colon. *Am J Surg* 2004; 187: 497-500.

Ward J, Robinson PJ, Guthrie JA et al. Liver Metastases in candidates for hepatic resection: comparison of helical CT and gadolinium- and SPIO-enhanced MR imaging. *Radiology* 2005; 237: 170-180.

Watabe H, Yamaji Y, Okamoto M, Ohta M, Ikenoue T et al. Risk assessment for delayed hemorrhagic complication of colonic polypectomy: Polyp-related factors and patient related factors. *Gastrointest Endosc* 2006 Jul; 64 (1): 73-8.

Watson Aj, Shanmugam V, Mackay I et al. Outcomes after placement of colorectal stents. *Colorectal Disease* 2005; 71 (1): 70-3.

Watson P, Vasen HF, Mecklin JP, Bernstein I, Aarnio M, Järvinen HJ et al. The risk of extra-colonic, extra-endometrial cancer in the Lynch syndrome. *Int J Cancer* 2008; 123: 444-449.

Watt Am, Faragher IG, Griffin TT, Rieger NA, Maddern GJ. Self-expanding metallic stents for relieving malignant colorectal obstruction: a systematic review. *Ann Surg* 2007 Jul; 246 (1): 24-30.

Wells BJ, Stotland P, Ko MA, Al-Susukhni W, Wunder J, Ferguson P et al. Results of an Aggressive Approach to Resection of Locally Recurrent Rectal Cancer. *Ann Surg Oncol* 2007; 14 (2): 390-95.

West NP, Morris EJ, Rotimi O, Cairns A, Finan PJ, Quirke P. Pathology grading of colon cancer surgical resection and its association with survival: a retrospective observational study. *Lancet Oncol* 2008; Jul 28.

Wexner SD, Cohen SM. Port site metastases after laparoscopic colorectal surgery for cure of malignancy. *Br J Surg* 1995; 82: 295-298.

White CM, Macfie J. Immediate colectomy and primary anastomosis for acute obstruction due to carcinoma of the left colon and rectum. *Dis Colon Rectum* 1985; 28: 155-7.

White P, Kehlet H (2007). Improving pain management: are we jumping from the frying pan into the fire? *Anesth Analg* 105: 10-12.

Whitehouse PA, Armitage JN, Tilney HS, Simson JN. Transanal endoscopic microsurgery: local recurrence rate following resection of rectal cancer. *Colorectal Dis* 2008; 10: 187-93.

WHO. Tumors of the digestive system. IARC 2000.

Whynes DK, Neilson AR, Robinson MH, Hardcastle JD. Colorectal cancer screening and quality of life. *Qual Life Res* 1994; 3: 191-8.

Whynes DK, Frew EJ, Edwards R, Atkin WS. Costs of flexible sigmoidoscopy screening for colorectal cancer in the United Kingdom. *Int J Technol Assess Health Care* 2003 Spring; 19: 384-95.

Wibe A, Syse A, Andersen E, Tretli S, Myrvold HE, Soreide O. Norwegian Rectal Cancer Group. Oncological outcomes after total mesorectal excision for cure for cancer of the lower rectum: Anterior vs. abdominoperineal resection. *Dis Colon Rectum* 2004; 47: 48-58.

Wibe A, Eriksen MT, Syse A et al. Effect of hospital caseload on long-term outcome after standardization of rectal cancer surgery at a national level. *Br J Surg* 2005; 92: 217-24.

Wibe A, Möller B, Norstein J et al. A national strategic change in treatment policy for rectal cancer-implementation of total mesorectal excision as routine treatment in Norway. A national audit. *Dis Colon Rectum* 2002; 45: 857-866.

Wibe A, Rendedal PR, Svensson E, Norstein J, Eide TJ, Myrvold HE et al. Prognostic significance of the circumferential resection margin following total mesorectal excision for rectal cancer. *Br J Surg* 2002;K 89: 327-34.

Wibe A. Magnetic resonance imaging for rectal cancer. *nat Clin Pract Oncol* 2007; 4: 222-3.

Wiering B, Ruers TJ, Krabbe PF, Dekker HM, Oyen WJ. Comparison of multiphase CT, FDG-PET and intra-operative ultrasound in patients with colorectal liver metastase selected for surgery. *Ann Surg Oncol* 2007; 14: 818-826.

Wiggers T, Jeekel J, Arends JW, Brinkhorst AP, Kluck HM, Luyk CI et al. No-touch isolation technique in colon cancer: A controlled prospective trial. *Br J Surg* 1988a; 75: 409-15.

Wiggers T, Arends JW, Volvics A. Regression analysis of prognostic factors in colorectal cancer after curative resection. *Dis Colon Rectum* 1988b; 31: 33-41.

Wiggers T, Mannaerts GHH, Marinelli AWKS, Martijn H, Rutten HJT. Surgery for Locally Recurrent Rectal Cancer. *Colorectal Dis* 2003; 5: 504-07.

Wiig JN, Dueland S, Larsen S, Giercksky K-E. Behandling av lokalt residiv ved endetarmskreft. *Tidsskr Nor Lægeforen* 2007; 127: 3097-101.

Wille-Jørgensen P, Guenaga KF, Castro AA, Matos D (2003). Clinical value of preoperative mechanical bowel cleansing in elective colorectal surgery: a systematic review. *Dis Colon Rectum* 46: 1013-1020.

Wille-Jørgensen P, Kjaergaard J, Jørgensen T, Korsgaard LT. Failure in prophylactic management of thromboembolic disease in colorectal surgery. *Dis Colon Rectum* 1988; 31: 384-386.

Wille-Jørgensen P, Rasmussen MS, Andersen BR, Borly L. Heparins and mechanical methods for thromboprophylaxis in colorectal surgery (Cochrane review). *The Cochrane Library*, issue 1. 2002. Oxford, Update Software.

Williams NS, Dixon MF, Johnston D. Reappraisal of the 5 centimetre rule of distal excision for carcinoma of the rectum: a study of distal intramural spread and of patients' survival. *Br J Surg* 1983; 70: 150-4.

Wilson RG, Gollock JM. Obstructing carcinoma of the left colon managed by subtotal colectomy. *J R Coll Surg Edinb* 1989; 34: 25-6.

Winawer S, Fletcher R, Rex D, Bond J, Burt R, Ferrucci J et al. Colorectal cancer screening and surveillance: clinical guidelines and rationale—update based on new evidence. *Gastroenterology* 2003; 124: 544-560.

Winawer SJ, Fletcher RH, Miller L, Godlee F, Stolar MH, Mulrow CD et al. Colorectal cancer screening: clinical guidelines and rationale. *Gastroenterology* 1997; 112: 594-642.

Winawer SJ, Zauber AG, Ho MN et al. Prevention of colorectal cancer by colonoscopic polypectomy. The National Polyp Study Workshop. *N Eng J Med* 1993; 329: 1977-1981.

Wind J, Polle S, Fung Kon Jin P, Dejon C, von Meyenfeldt M, Ubbink D et al (2006). Systematic review of enhanced recovery programmes in colonic surgery. *Br J Surg* 93: 800-809.

Winde G, Nottberg H, Keller R, Schmid Kw, Bunte H. Surgical cure for early rectal carcinomas (T1). Transanal endoscopic microsurgery vs. anterior resection. *Dis Colon Rectum* 1996; 39: 969-76.

Wolin KY, Lee IM, Colditz GA, Glynn RJ, Fuchs C, Giovannucci E. Leisure-time physical activity patterns and risk of colon cancer in women. 1: *Int J Cancer* 2007; 121: 2776-81.

Wolmark N, Fisher B, Rockette H, Redmond C, Wickerham DL, Fisher ER et al. "Postoperative adjuvant chemotherapy or BCG for colon cancer: results from NSABP protocol C-01". *J Natl Cancer Inst*, Vol. 80, No. 1, 1998, pp. 30-36.

Wong PF, Gilliam AD, Kumar S, Shenfine J, O'Dair GN, Leaper DJ. 2005. Antibiotic regimens for secondary peritonitis of gastrointestinal origin in adults. *Cochrane Database. Syst.Rev.*:CD004539.

Wong RKS, Tanda V, De Silva S, Figueredo A. Pre-operative radiotherapy and curative surgery for the management of localized rectal carcinoma. *Cochrane Database of Systematic Reviews* 2007, Issue 2. Art. No.:L CD002102.

Wu GH, Liu ZH, Wu ZG (2006). Perioperative artificial nutrition in malnourished gastrointestinal cancer patients. *World J Gastroenterol* 12: 2441-2444.

Yamada K, Ishizawa T, Niwa K, Chuman Y, Akiba S, Aikou T. Locally Recurrent Rectal Cancer. *Br J Surg* 2001; 88: 988-93.

Ye F, Lin JJ. Clinical application of biofragmental anastomosis ring for intestinal anastomosis. *Zhejiang Da Xue Xue Bao Ti Xue Ban*. 2006 Nov; 356 (6): 668-72.

Young GP, St John DJ, Winawer SJ, Rozen P, WHO (World Health Organization) and OMED (World Organization for Digestive Endoscopy). Choice of fecal occult blood test for colorectal cancer screening: recommendations based on performance characteristics in population studies: a WHO (World Health Organization) and OMED (World Organization for Digestive Endoscopy) report. *Am J Gastroenterol* 2002; 97: 2499-507.

Zacharakis E, Freilich S, Rekhraj S, Athanasiou T, Paraskeva P, Ziprin P et al. Transanal endoscopic microsurgery for rectal tumors: the St. Mary's experience. *Am J Surg* 2007; 194: 694-8.

Zerz A, Müller-Stich BP, Beck J, Linke GR, Tarantino I, Lange J. Endoscopic posterior mesorectal resection after transanal local excision of T1 carcinomas of the lower third of the rectum. *Dis Colon Rectum* 2006; 49: 919-24.

Zhou ZG, Hu M, Li Y, Lei WZ, Cheng Z et al. Laparoscopic versus open total mesorectal excision with anal sphincter preservation for low rectal cancer. *Surg Endosc* 2004; 18: 1211-1215.

Zirngibl H, Husemann B, Hermanek P. Intraoperative spillage of tumour cells in surgery for rectal cancer. *Dis Colon Rectum* 1990; 33: 610-4.

Zlobec I, Baker K, Minoo P et al. Node-Negative Colorectal Cancer at High Risk of Distant Metastasis Identified by Combined Analysis of Lymph Node Status, Vascular Invasion and Raf-1 Kinase Inhibitor Prein Expression. *Clin Cancer Res* 2008; 14 (1): 143-148.

Zmora O, Gervaz P, Wexner SD. Trocar site recurrence in laparoscopic surgery for colorectal cancer. *Surg Endosc* 2001; 15: 788-793.

Zmora O, Lebedyev A, Hoffman A, Khaikin M, Munz Y, Shabtai M et al. Laparoscopic colectomy without mechanical bowel preparation. *Int J Colorectal Dis* 2006; 21: 683-687.

Zmora O, Mahajna A, Bar-Zakai B, Rosin D, Hershko D, Shabtai M et al (2003). Colon and rectal surgery without mechanical bowel preparation: a randomized prospective trial. *Ann Surg* 237: 363-367.

Öhman U. Prognosis in patients with obstructing colorectal carcinoma. *Am J Surg* 1982; 143: 742-7.

Årsrapport 2006. Landsdækkende database for kræft i tyktarm og endetarm. Dansk Kolorektalcancer Database, 2007.